

NONCOMMUTATIVE PROJECTIVE CURVES AND QUANTUM LOOP ALGEBRAS

OLIVIER SCHIFFMANN

ABSTRACT. We show that the Hall algebra of the category of coherent sheaves on a weighted projective line over a finite field provides a realization of the (quantized) enveloping algebra of a certain nilpotent subalgebra of the affinization of the corresponding Kac-Moody algebra. In particular, this yields a geometric realization of the quantized enveloping algebra of elliptic (or 2-toroidal) algebras of types $D_4^{(1,1)}$, $E_6^{(1,1)}$, $E_7^{(1,1)}$ and $E_8^{(1,1)}$ in terms of coherent sheaves on weighted projective lines of genus one, or equivalently in terms of equivariant coherent sheaves on elliptic curves.

Dedicated to Igor Frenkel on the occasion of his 50th birthday

Introduction. The geometric approach to quantum groups developed in the past 15 years is based on a deep relationship between simple or affine (and to a lesser extent Kac-Moody) Lie algebras and certain finite-dimensional hereditary algebras. More precisely, let $\mathfrak{g}$ be a Kac-Moody algebra and Γ its Dynkin diagram. Ringel proved in [Ri1] that the Hall algebra of the category of representations, over a finite field $\mathbb{F}_q$, of a quiver whose underlying graph is Γ provides a realization of the “positive part” $\mathbf{U}_q^+(\mathfrak{g})$ of the Drinfeld-Jimbo quantum group associated to $\mathfrak{g}$. This result was the starting point of Lusztig’s geometric construction of the canonical basis of $\mathbf{U}_q^+(\mathfrak{g})$ (see [Lu]).

Another natural example of a hereditary category is provided by the category $\text{Coh}(X)$ of coherent sheaves on a smooth projective curve X . In the remarkable paper [Kap], Kapranov observed a striking analogy between a function field analog of the algebra of unramified automorphic forms (for $GL(N)$ for all N) and Drinfeld’s loop-like realization of quantum affine algebras. In particular, his result provides an isomorphism between a natural subalgebra of the Hall algebra of the category $\text{Coh}(\mathbb{P}^1(\mathbb{F}_q))$ and Drinfeld’s “positive part” $\mathbf{U}_q^+(\widehat{\mathfrak{sl}}_2)$.

In this paper we extend Kapranov’s result. Rather than considering higher genus smooth projective curves (whose Hall algebras are wild) we study the Hall algebra of the category of coherent sheaves on certain “noncommutative smooth projective curves”- the so-called *weighted projective lines*, introduced by Geigle and Lenzing ([GL]). These fundamental examples of “noncommutative smooth projective curves” have attracted some attention lately (see e.g [Hap2] and [CB]). Our main result (Theorem 5.2) provides, when $\mathfrak{g}$ is a simple Lie algebra or an affine Lie algebra of type $D_4^{(1)}$, $E_6^{(1)}$, $E_7^{(1)}$ or $E_8^{(1)}$, a natural embedding $\mathbf{U}_q(\widehat{\mathfrak{n}}) \hookrightarrow \mathbf{H}_{\text{Coh}(\mathbb{X}_{\mathfrak{g}})}$ of a certain “positive part” of $\mathbf{U}_q(\mathfrak{L}\mathfrak{g})$ into the Hall algebra of a suitable weighted projective line $\mathbb{X}_{\mathfrak{g}}$. Here $\mathfrak{L}\mathfrak{g}$ is the universal central extension of $\mathfrak{g}[t, t^{-1}]$.

In particular, this gives a geometric construction of the quantum toroidal algebras of type D_4, E_6, E_7 and E_8 . In this case, the two loops have a natural interpretation as the two discrete invariants (rank and degree) of a coherent sheaf on $\mathbb{X}_{\mathfrak{g}}$. Note that the exceptional role played by these four types among all affine Dynkin diagrams is well-known in the theory of quadratic forms (see e.g [Ri3]). Our

computations extend those of Bauman and Kassel in [BK], where the case of the projective line $\mathbb{P}^1$ is considered in details. Our proof also makes use in an essential way of the Harder-Narasimhan filtration on semistable coherent sheaves and on the theory of mutations. The use of the Harder-Narasimhan filtration in the context of Hall algebras is also noted in [Re].

Recently, Y. Lin and L. Peng independently obtained another construction of the enveloping algebra of double-loop algebras of type D_4, E_6, E_7 and E_8 by considering the root category (as in [PX]) of certain tame algebras $\Lambda_{\mathfrak{g}}$ (the *tubular algebras*). Our construction is related to theirs by the existence, in this case, of a derived equivalence

$$D^b(\text{Coh}(\mathbb{X}_{\mathfrak{g}})) \simeq D^b(\text{Rep}(\Lambda_{\mathfrak{g}})).$$

Finally, we note the following link of our work with the McKay correspondence. Let $\mathfrak{g}$ be a simple Lie algebra of type A_{2n}, D_n or E_k for $k = 6, 7, 8$, and let $\Gamma \supset \{\pm 1\}$ be the finite subgroup of $SL(2, \mathbb{C})$ associated to $\mathfrak{g}$ by the McKay correspondence, and set $\Gamma' = \Gamma/\{\pm 1\} \in SO(3)$. When the base field is algebraically closed, the (ramified) quotient $\mathbb{P}^1/\Gamma'$ and the (smooth, noncommutative) curve $\mathbb{X}_{\mathfrak{g}}$ are related by an equivalence of categories $\text{Coh}(\mathbb{X}_{\mathfrak{g}}) \simeq \text{Coh}_{\Gamma'}(\mathbb{P}^1)$ and a derived equivalence

$$D^b(\text{Coh}(\mathbb{X}_{\mathfrak{g}})) \simeq D^b(\Lambda_{\mathfrak{g}}) \simeq D_{\Gamma'}^b(\text{Coh}(\mathbb{P}^1)),$$

where $\Lambda_{\mathfrak{g}}$ is the path algebra of the McKay quiver of Γ , and $D_{\Gamma'}^b(\text{Coh}(\mathbb{P}^1))$ is the Γ' -equivariant derived category of $\text{Coh}(\mathbb{P}^1)$. Furthermore, the minimal desingularization $\widetilde{T^*(\mathbb{P}^1)/\Gamma'}$ is isomorphic to the minimal desingularization $\widetilde{\mathbb{C}^2/\Gamma}$ of the Kleinian singularity $\mathbb{C}^2/\Gamma$ (see [La]). Such spaces (the so-called ALE spaces), and more generally the moduli space of vector bundles on them, have recently been used by Nakajima in his geometric construction of representations of $\widehat{\mathfrak{g}}$ (see [N]).

Now let $\mathfrak{g}$ be of type $D_4^{(1)}, E_6^{(1)}, E_7^{(1)}$ or $E_8^{(1)}$. As shown in [GL], §5.8, we now have $\text{Coh}(\mathbb{X}_{\mathfrak{g}}) \simeq \text{Coh}_G(\mathcal{E})$ where $\mathcal{E}$ is an elliptic curve and G a finite subgroup of $\text{Aut}(\mathcal{E})$. This suggests the study of the “noncommutative space” $T^*\mathbb{X}_{\mathfrak{g}}$ and the moduli space of vector bundles on it in the spirit of Nakajima’s work.

More generally, as explained to us by H. Lenzing, there exists, for *any* weighted projective line $\mathbb{X}_{\mathfrak{g}}$ with at least three marked points, a pair (X, G) consisting of a smooth projective curve X and a finite group of automorphisms $G \subset \text{Aut}(X)$ such that $\text{Coh}_G(X) \simeq \text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\Delta}})$. Conversely, if X is a smooth projective curve and G a group of automorphisms of X such that $X/G \simeq \mathbb{P}^1$ then $\text{Coh}_G(X)$ is equivalent to the category of coherent sheaves on the weighted projective line $\mathbb{X}_{\mathbf{p}, \underline{\Delta}}$, where $\underline{\Delta}$ and $\mathbf{p}$ are the ramification locus and multiplicities respectively.

Acknowledgements.

I would like to thank John Duncan, Igor Frenkel, Michael Kapranov and Markus Reineke for interesting discussions, as well as B. Enriquez, L. Hille, D. Kussin, H. Lenzing and C. Ringel. I am grateful to Hadi Salmasian for teaching me all I know about Farey sequences.

1. LOOP ALGEBRAS AND QUANTUM GROUPS

1.1. Let $A = (a_{ij})_{i,j=1}^r$ be an irreducible symmetric generalized Cartan matrix of simply laced type and let $(\mathfrak{h}, \Pi, \Pi^\vee)$ be a realization of A (see [Kac]), where $\Pi = \{\alpha_1, \dots, \alpha_r\} \subset \mathfrak{h}^*$ and $\Pi^\vee = \{\alpha_1^\vee, \dots, \alpha_r^\vee\} \subset \mathfrak{h}$. Let $\mathfrak{h}' = \text{Span}_{\mathbb{C}}\{\alpha_1^\vee, \dots, \alpha_r^\vee\}$

and let $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h}' \oplus \mathfrak{n}_+$ be the associated complex Kac-Moody algebra, with generators $\{e_i, f_i \mid i = 1, \dots, r\} \cup \mathfrak{h}'$ satisfying the following set of relations

$$\begin{aligned} [\mathfrak{h}', \mathfrak{h}'] &= 0, & [e_i, f_j] &= \delta_{ij} \alpha_i^\vee, \\ [\alpha_j^\vee, e_i] &= a_{ij} e_{\alpha_i}, & [\alpha_j^\vee, f_i] &= -a_{ij} f_i, \\ ad(e_i)^{1-a_{ij}}(e_j) &= 0, & ad(f_i)^{1-a_{ij}}(f_j) &= 0. \end{aligned}$$

Let $(,)$ be a standard invariant bilinear form on $\mathfrak{g}$ (see [Kac]). In particular, we have $(e_i, f_i) = 1$ for $i = 1, \dots, r$. Let us further denote by $\Delta \subset \mathfrak{h}^*$ the root system of $\mathfrak{g}$ and by $Q = \bigoplus_i \mathbb{Z} \alpha_i$ the root lattice. We also let $\Delta^{im} \subset \Delta$ stand for the subset of imaginary roots. The $\mathbb{Z}$ -module Q is equipped with the Cartan bilinear form determined by the relation $(\alpha_i, \alpha_j) = a_{ij}$.

1.2. Let $\widehat{\mathfrak{g}} = \mathfrak{g}[t, t^{-1}] \oplus \mathbb{C}\mathbf{c}$ be the affinization of $\mathfrak{g}$, where $\mathbf{c}$ is a central element, and where the Lie bracket is given by

$$[xt^n, yt^m] = [x, y]t^{n+m} + n\delta_{n,-m}(x, y)\mathbf{c}.$$

The algebra $\widehat{\mathfrak{g}}$ is naturally equipped with a $\widehat{Q} = Q \oplus \mathbb{Z}\delta$ -grading, where

$$\widehat{\mathfrak{g}}[\alpha + l\delta] = \mathfrak{g}[\alpha]t^l, \quad \widehat{\mathfrak{g}}[0] = \mathfrak{h} \oplus \mathbb{C}\mathbf{c}, \quad \widehat{\mathfrak{g}}[l\delta] = \mathfrak{h}t^l.$$

We extend the Cartan form to $\widehat{Q}$ by setting $(\delta, \alpha) = 0$ for all $\alpha \in \widehat{Q}$. The root system of $\widehat{\mathfrak{g}}$ is $\widehat{\Delta} = \mathbb{Z}^*\delta \cup \{\Delta + \mathbb{Z}\delta\}$. We will say that a root $\alpha \in \widehat{\Delta}$ is *real* if $(\alpha, \alpha) = 2$ and *imaginary* if $(\alpha, \alpha) \leq 0$.

1.3. We will consider certain extensions of the Lie algebra $\widehat{\mathfrak{g}}$. Let us denote by $\mathfrak{Lg}$ the Lie algebra generated by elements $h_{i,k}, e_{i,k}, f_{i,k}$ for $i = 1, \dots, r$, $k \in \mathbb{Z}$ and $\mathbf{c}$ subject to the following relations :

$$\begin{aligned} [h_{i,k}, h_{j,l}] &= k\delta_{k,-l}a_{ij}\mathbf{c}, & [e_{i,k}, f_{j,l}] &= \delta_{i,j}h_{i,k+l} + k\delta_{k,-l}\mathbf{c}, \\ [h_{i,k}, e_{j,l}] &= a_{ij}e_{j,l+k}, & [h_{i,k}, f_{j,l}] &= -a_{ij}f_{j,k+l}, \\ [e_{i,k+1}, e_{j,l}] &= [e_{i,k}, e_{j,l+1}], & [f_{i,k+1}, f_{j,l}] &= [f_{i,k}, f_{j,l+1}] \\ [e_{i,k_1}, [e_{i,k_2}, [\dots [e_{i,k_n}, e_{j,l}] \dots]] &= 0 & \text{if } n = 1 - a_{ij}, \\ [f_{i,k_1}, [f_{i,k_2}, [\dots [f_{i,k_n}, f_{j,l}] \dots]] &= 0 & \text{if } n = 1 - a_{ij}. \end{aligned}$$

It is clear that $e_i \mapsto e_{i,0}, f_i \mapsto f_{i,0}, \alpha_i^\vee \mapsto h_{i,0}$ defines an embedding $\mathfrak{g} \subset \mathfrak{Lg}$. Moreover, giving generators $e_{i,k}, f_{i,k}, h_{i,k}$ degrees $\alpha_i + k\delta$, $-\alpha_i + k\delta$ and $k\delta$ respectively endows $\mathfrak{Lg}$ with a $\widehat{Q}$ -grading.

There is a surjective Lie algebra morphism $\varphi : \mathfrak{Lg} \rightarrow \widehat{\mathfrak{g}}$ given by $e_{i,k} \mapsto e_i t^k$, $f_{i,k} \mapsto f_i t^k$, $h_{i,k} \mapsto \alpha_i^\vee t^k$, $\mathbf{c} \mapsto \mathbf{c}$. It is proved in [MRY] that

$$Ker \varphi \subset \bigoplus_{\substack{\alpha \in \Delta^{im} \\ k \in \mathbb{Z}}} \mathfrak{Lg}[\alpha + k\delta].$$

In particular, the root system of $\mathfrak{Lg}$ is also $\widehat{\Delta}$ and the weight spaces corresponding to real roots are one-dimensional.


When $\mathfrak{g}$ is a simple complex Lie algebra, it is known that φ is an isomorphism $\mathfrak{L}\mathfrak{g} \simeq \widehat{\mathfrak{g}}$. Now suppose that $\mathfrak{g} = \mathfrak{g}_0[s, s^{-1}] \oplus \mathbb{C}\mathbf{c}_0 = \widehat{\mathfrak{g}}_0$ is an (untwisted) affine algebra. According to [MRY], $\mathfrak{L}\mathfrak{g}$ is then the universal central extension of $\mathfrak{g}_0[s, s^{-1}, t, t^{-1}]$ (the toroidal algebra, or elliptic algebra). Let $\delta_0 \in \Delta$ stand for the imaginary root of $\widehat{\mathfrak{g}}_0$, and let $\Delta_0 \subset \Delta$ be the root system of $\mathfrak{g}_0$, so that $\Delta = \mathbb{Z}^*\delta_0 \cup (\Delta_0 + \mathbb{Z}\delta_0)$, and

$$\widehat{\Delta} = (\mathbb{Z}\delta_0 + \mathbb{Z}\delta) \cup (\Delta_0 + \mathbb{Z}\delta_0 + \mathbb{Z}\delta) \setminus \{0\}.$$

We have (see [MRY], Proposition 3.6)

$$(1.1) \quad \dim \mathfrak{L}\mathfrak{g}[\alpha + k_0\delta_0 + k\delta] = \begin{cases} 1 & \text{if } \alpha \neq 0 \\ l+1 & \text{if } \alpha = 0, (k_0, k) \neq (0, 0) \\ l+2 & \text{if } \alpha = k_0 = k = 0 \end{cases}$$

1.4. We will now consider certain nilpotent subalgebras in $\mathfrak{L}\mathfrak{g}$ and $\widehat{\mathfrak{g}}$. Let $p_1, \dots, p_n \in \mathbb{N}$ and let us consider the “star” Dynkin diagram



Let us denote by $\star$ the central node and let $J_1, \dots, J_n$ be the branches, which are subdiagrams are of type $A_{p_1-1}, \dots, A_{p_n-1}$ respectively. We can assume that $p_i > 1$ for all i . These examples include all finite type Dynkin diagrams as well as the affine Dynkin diagrams of types $D_4^{(1)}, E_6^{(1)}, E_7^{(1)}$, and $E_8^{(1)}$.

Let $\mathfrak{g}$ be the Kac-Moody algebra corresponding to Γ , and let $\mathfrak{p}$ be the maximal parabolic subalgebra of $\mathfrak{g}$ associated to the node $\star$. Denote by $\mathfrak{l}$ and $\mathfrak{m}$ its standard Levi and nilpotent radical respectively. Let us write $\mathfrak{l} = \mathfrak{n}_\mathfrak{l}^+ \oplus \mathfrak{h} \oplus \mathfrak{n}_\mathfrak{l}^-$ for the standard Cartan decomposition of $\mathfrak{l}$ and set

$$\widehat{\mathfrak{l}} := \mathfrak{n}_\mathfrak{l}^+ \oplus t\mathfrak{l}[t] \subset \mathfrak{l}[t, t^{-1}], \quad \widehat{\mathfrak{f}} := \mathfrak{m}[t, t^{-1}].$$

Finally, we put

$$(1.2) \quad \widehat{\mathfrak{n}}' := \widehat{\mathfrak{f}} \oplus \widehat{\mathfrak{l}} \subset \widehat{\mathfrak{g}}.$$

The following description of $\widehat{\mathfrak{n}}'$ will be convenient for us. For every $s = 1, \dots, n$ let $\mathfrak{g}^s$ be the subalgebra of $\mathfrak{g}$ generated by the elements $\{e_j, f_j, \alpha_j^\vee \mid j \in J_s\}$. Let θ_s be the highest root of $\mathfrak{g}^s$ and let $f_{\theta_s} \in \mathfrak{g}_{-\theta_s}^s$ be any lowest root vector. Let $\widehat{\mathfrak{n}}^s \subset \widehat{\mathfrak{g}}$ be generated by $\{e_j \mid j \in J_s\} \cup \{f_{\theta_s}t\}$. Then $\widehat{\mathfrak{n}}'$ is the subalgebra generated by

$$(1.3) \quad \bigcup_{s=1}^n \widehat{\mathfrak{n}}^s \cup \{e_\star t^l \mid l \in \mathbb{Z}\} \cup \{h_\star t^r \mid r \in \mathbb{N}^*\}.$$

and $\widehat{\mathfrak{l}}$ is the subalgebra generated by $\widehat{\mathfrak{n}}^s$ for $s = 1, \dots, n$ and $h_\star t^r$ for $r \geq 1$.

Observe that, for any i , the subalgebra of $\mathfrak{L}\mathfrak{g}$ generated by $e_{j,l}, f_{j,l}, h_{j,l}$ for $j \in J_i$ and $l \in \mathbb{Z}$ is canonically isomorphic (via the map φ) to an affine algebra of type A_{p_i-1} , and we can define the subalgebra $\widehat{\mathfrak{n}}^s \subset \mathfrak{L}\mathfrak{g}$ in the same way as above. We now define $\mathfrak{L}\mathfrak{n}$ to be the subalgebra of $\mathfrak{L}\mathfrak{g}$ generated by

$$(1.4) \quad \bigcup_{s=1}^n \widehat{\mathfrak{n}}^s \cup \{e_{\star,l} \mid l \in \mathbb{Z}\} \cup \{h_{\star,r} \mid r \in \mathbb{N}^*\}.$$

1.5. Let $p > 0$. The Lie algebra $\mathfrak{gl}_p$ of p by p matrices has a natural basis

$$\{E_{ij} \mid i, j = 1, \dots, p\}$$

such that $[E_{ij}, E_{kl}] = \delta_{jk}E_{il} - \delta_{il}E_{kj}$. Set $h_0 = -E_{11}$. Then

$$\mathfrak{gl}_p = \mathfrak{sl}_p \oplus \mathbb{C}h_0.$$

Let $(x, y) = \text{Tr}(xy)$ be the (normalized) Killing form. Define $\widehat{\mathfrak{gl}}_p = \mathfrak{gl}_p[t, t^{-1}] \oplus \mathbb{C}c$ as in Section 1.2. Set $e_i = E_{i,i+1}$ for $i = 1, \dots, p-1$ and $e_0 = E_{p1} \otimes t$. Let $\widehat{\mathfrak{sl}}_p^+ \subset \widehat{\mathfrak{gl}}_p$ (resp. $\widehat{\mathfrak{gl}}_p^+ \subset \widehat{\mathfrak{gl}}_p$) be the subalgebra generated by e_i , $i = 0, \dots, p-1$ (resp. generated by e_i , $i = 1, \dots, p-1$ and $\mathfrak{gl}_p \otimes t\mathbb{C}[t]$). In particular,

$$(1.5) \quad \mathbf{U}(\widehat{\mathfrak{gl}}_p^+) = \mathbf{U}(\widehat{\mathfrak{sl}}_p^+) \otimes \mathbb{C}[h_1, h_2, \dots],$$

where we set $h_i = h_0 t^i$.

1.6. Let Γ be as in Section 1.4. We give a presentation of (a certain extension of) the Lie algebra $\widehat{\mathfrak{n}}'$ by generators and relations. For each s we let $\phi_s : A_{p_s-1} \xrightarrow{\sim} J_s$ be the unique bijection such that $\phi_s(1)$ is adjacent to the central node $\star$.

Definition. Let $\widehat{\mathfrak{n}}$ be the Lie algebra generated by ε_j for $j \in \Gamma \setminus \{\star\}$, $\varepsilon_0^{(s)}$ for $s = 1, \dots, n$, and $\varepsilon_{\star,t}$, $\theta_{\star,r}$, for $r \geq 1$ and $t \in \mathbb{Z}$, subject to the following set of relations :

- i) For all s , the assignment $e_j \mapsto \varepsilon_{\phi_s(j)}$, $e_0 \mapsto \varepsilon_0^{(s)}$ and $h_r \mapsto \theta_{\star,r}$ extends to an embedding $\widehat{\mathfrak{gl}}_{p_s}^+ \hookrightarrow \widehat{\mathfrak{n}}$, and $[\widehat{\mathfrak{gl}}_{p_s}^+, \widehat{\mathfrak{gl}}_{p_{s'}}^+] = 0$ if $s \neq s'$.
- ii) We have

$$(1.6) \quad [\theta_{\star,r}, \varepsilon_{\star,t}] = 2\varepsilon_{\star,r+t}, \quad [\varepsilon_{\star,t}, \varepsilon_{\star,t'}] = 0$$

iii)

$$(1.7) \quad a_{\star j} = 0 \Rightarrow [\varepsilon_{\star,t}, \varepsilon_j] = 0, \quad [\varepsilon_{\star,t}, \varepsilon_0^{(s)}] = 0, \quad s = 1, \dots, n$$

$$(1.8) \quad p_s > 2 \Rightarrow [\varepsilon_0^{(s)}, [\varepsilon_{\star,t}, \varepsilon_{\phi_s(1)}]] = 0.$$

- iv) For $s = 1, \dots, n$ and $j = \phi_s(1)$ set $\varepsilon_{j,r} = [\varepsilon_j, \theta_{\star,r}]$. Then for any $t, t_1, t_2 \in \mathbb{Z}$ and $r, r_1, r_2 \geq 1$,

$$(1.9) \quad [\varepsilon_{\star,t_1}, [\varepsilon_{\star,t_2}, \varepsilon_{j,r}]] = [\varepsilon_{j,r_1}, [\varepsilon_{j,r_2}, \varepsilon_{\star,t}]] = 0$$

$$(1.10) \quad [\varepsilon_{\star,t}, \varepsilon_{j,r+1}] = [\varepsilon_{\star,t+1}, \varepsilon_{j,r}]$$

It is easy to check that the assignment $\varepsilon_{\star,l} \mapsto e_{\star,l}$, $\theta_{\star,k} \mapsto h_{\star,k}$, $\varepsilon_j \mapsto e_j$, $\varepsilon_0^{(s)} \mapsto f_{\theta_s} t$ extends to a surjective homomorphism $\phi : \widehat{\mathfrak{n}} \rightarrow \mathfrak{L}\mathfrak{n}$. Moreover, $\widehat{\mathfrak{n}}$ is $\widehat{Q}$ -graded.

Proposition 1.1. *We have*

$$\text{Ker } \phi \subset \bigoplus_{\substack{\alpha \in \Delta^{im} \\ l \in \mathbb{Z}}} \widehat{\mathfrak{n}}[\alpha + l\delta]$$

Moreover, ϕ is an isomorphism if Γ is a finite or an affine Dynkin diagram.

Proof. For $s = 1, \dots, n$ we denote by $\mathfrak{n}_-^s \subset \mathfrak{sl}_{p_s}$ the subspace spanned by $E_{k,l}$ for $k > l$.

Lemma 1.1. *For any s , any $t \in \mathbb{Z}$ and any $x \otimes t^r \in \mathfrak{n}_-^s \otimes t\mathbb{C}[t] \subset \widehat{\mathfrak{gl}}_{p_s}^+$ we have $[\varepsilon_{\star,t}, x \otimes t^r] = 0$.*

Proof. Set $\varepsilon_{0,r}^s = [\theta_{\star,r}, \varepsilon_0^{(s)}]$. If $p_s = 2$ then the lemma follows from (1.7). If $p_s > 2$, observe that $[\varepsilon_{0,r}, \varepsilon_{\star,t}] = 0$ and that

$$\begin{aligned}
 (1.11) \quad [\varepsilon_{\star,t}, [\varepsilon_{0,r}, \varepsilon_1]] &= [\varepsilon_{0,r}, [\varepsilon_{\star,t}, \varepsilon_1]] \\
 &= [[\theta_{\star,r}, \varepsilon_0], [\varepsilon_{\star,t}, \varepsilon_1]] \\
 &= [\theta_{\star,r}, [\varepsilon_0, [\varepsilon_{\star,t}, \varepsilon_1]]] + [\varepsilon_0, [\theta_{\star,r}, [\varepsilon_{\star,t}, \varepsilon_1]]] \\
 &= 2[\varepsilon_0, [\varepsilon_{\star,t+r}, \varepsilon_1]] - [\varepsilon_0, [\varepsilon_{\star,t}, \varepsilon_{1,r}]] \\
 &= 0
 \end{aligned}$$

where we have used (1.8) and (1.10). Any $x \otimes t^r \in \mathfrak{n}_-^s \otimes t\mathbb{C}[t]$ is a linear combination of terms of the form $[\varepsilon_{0,r}, [\varepsilon_{i_1}, [\varepsilon_{i_2}, \dots]]]$ where $i_1, i_2, \dots \in \{1, \dots, p_s - 1\}$ are distinct. The lemma is now a consequence of (1.11). $\diamond$

The algebra $\widehat{\mathfrak{n}}$ is linearly spanned by iterated commutators

$$[\alpha_1, \dots, \alpha_r] = [\alpha_1, [\alpha_2, [\dots, \alpha_r] \dots]]$$

with $\alpha_i \in \{e_{\star,l}, e_j t^k, h_j t^k, x t^r \mid x \in \mathfrak{n}_-^s; s = 1, \dots, n; l \in \mathbb{Z}; k \geq 0; r \geq 1\}$. Set

$$\widehat{\mathfrak{n}}_{\pm} = \bigoplus_{\alpha \in \Delta^{\pm}, l \in \mathbb{Z}} \widehat{\mathfrak{n}}[\alpha + l\delta], \quad (\mathfrak{L}\mathfrak{n})_{\pm} = \bigoplus_{\alpha \in \Delta^{\pm}, l \in \mathbb{Z}} \mathfrak{L}\mathfrak{n}[\alpha + l\delta].$$

From the defining relations and the Lemma above one easily deduces that $\widehat{\mathfrak{n}}_+$ is generated by $\varepsilon_{\star,l}, \varepsilon_j t^k$ for $l \in \mathbb{Z}, j \in \Gamma \setminus \{\star\}$ and $k \geq 0$. Similarly, $\widehat{\mathfrak{n}}_-$ is generated by $\{x t^r \mid x \in \mathfrak{n}_-^s, r \geq 1, s = 1, \dots, n\}$. It is clear that $\text{Ker } \phi|_{\widehat{\mathfrak{n}}_-} = 0$. On the other hand, define $\widehat{\mathfrak{n}}_{E,+}$ to be the Lie algebra generated by elements $\varepsilon_{\star,l}, \varepsilon_{j,k}$ for $l \in \mathbb{Z}, j \in \Gamma \setminus \{\star\}$ and $k \geq 0$ subject to the relations

$$\begin{aligned}
 [\varepsilon_{i,k+1}, \varepsilon_{j,l}] &= [\varepsilon_{i,k}, \varepsilon_{j,l+1}] \quad \forall k, j, \forall i, j \in \Gamma, \\
 [\varepsilon_{i,k_1}, [\varepsilon_{i,k_2}, [\dots [\varepsilon_{i,k_n}, \varepsilon_{j,l}] \dots]]] &= 0 \quad \text{if } n = 1 - a_{ij}, \forall k_1, \dots, k_n, l.
 \end{aligned}$$

There is an obvious surjective map $\psi : \widehat{\mathfrak{n}}_{E,+} \rightarrow \widehat{\mathfrak{n}}_+$. But from [E], Proposition 1.5 it follows that

$$\text{Ker } \phi\psi \subset \bigoplus_{\alpha \in \Delta^{im}, l \in \mathbb{Z}} \widehat{\mathfrak{n}}_{E,+}[\alpha + l\delta],$$

and from [E], Proposition 1.6 we deduce that $\phi\psi$ is injective when Γ is finite or affine. This proves the Proposition. $\diamond$

Observe that if Γ is finite then $\widehat{\mathfrak{n}} = \mathfrak{L}\mathfrak{n} = \widehat{\mathfrak{n}}'$. Let us now assume that Γ is affine. Let $\widehat{\mathfrak{t}} \subset \widehat{\mathfrak{n}}$ be the subalgebra generated by $\widehat{\mathfrak{gl}}_{p_i}^+$ for $i = 1, \dots, n$, and let $\widehat{\mathfrak{f}} \subset \widehat{\mathfrak{n}}$ be its (unique) $\widehat{Q}$ -graded complement. It is clear that $\widehat{\mathfrak{t}} \simeq \widehat{\mathfrak{t}}'$. We will say that a root $\alpha = \sum_{i \in \Gamma} \lambda_i \alpha_i$ is $\star$ -positive if $\lambda_{\star} > 0$.

Corollary 1.1. *Assume that Γ is affine. Then*

$$(1.12) \quad \dim \widehat{\mathfrak{f}}[\alpha + k_0 \delta_0 + k\delta] = \begin{cases} 1 & \text{if } \alpha \in \Delta_0^+ \text{ and } \alpha + k_0 \delta_0 \text{ is } \star\text{-positive} \\ l + 1 & \text{if } \alpha = 0, k_0 > 0 \\ 0 & \text{otherwise} \end{cases}$$

where $l = |\Gamma| - 1$ is the rank of $\mathfrak{g}_0$.

1.7. For any $n \in \mathbb{N}$ we set

$$[n] = \frac{v^n - v^{-n}}{v - v^{-1}}, \quad [n]! = \prod_{i=1}^n [i]!, \quad \begin{bmatrix} n \\ m \end{bmatrix} = \frac{[n]!}{[n-m]![m]},$$

and put $\mathbb{C}_v = \mathbb{C}[v, v^{-1}]$. Let $\mathbf{U}$ be the $\mathbb{C}(v)$ -algebra generated by $\{E_i, F_i, K_i, K_i^{-1} \mid i = 1, \dots, r\}$ satisfying the following relations

$$\begin{aligned} K_i K_i^{-1} &= K_i^{-1} K_i = 1, & [K_i, K_j] &= 0, \\ K_i E_j K_i^{-1} &= v^{a_{ij}} E_j, & K_i F_j K_i^{-1} &= v^{-a_{ij}} F_j \\ E_i F_j - F_j E_i &= \delta_{ij} \frac{K_i - K_i^{-1}}{v - v^{-1}}, \\ \sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix} E_i^{1-a_{ij}-k} E_j E_i^k &= 0, & i \neq j, \\ \sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix} F_i^{1-a_{ij}-k} F_j F_i^k &= 0, & i \neq j. \end{aligned}$$

The quantum enveloping algebra $\mathbf{U}_v(\mathfrak{g})$ is by definition the $\mathbb{C}_v$ -subalgebra of $\mathbf{U}$ generated by K_i, K_i^{-1} for $i = 1, \dots, r$ and the divided powers

$$E_i^{(n)} := \frac{E_i^n}{[n]!}, \quad F_i^{(n)} := \frac{F_i^n}{[n]!}$$

for $i = 1, \dots, r$ and $n \geq 1$. As usual, we let $\mathbf{U}_v^+(\mathfrak{g})$ be the $\mathbb{C}_v$ -subalgebra of $\mathbf{U}_v(\mathfrak{g})$ generated by $E_i^{(n)}$ for $i = 1, \dots, r$ and $n \geq 1$.

1.8. The quantum loop algebra (with zero central charge) $\mathbf{U}_v(L\mathfrak{g})$ is the $\mathbb{C}(v)$ -algebra generated by $x_{i,k}^\pm, h_{i,l}$ and $K_i^\pm$ for $i = 1, \dots, r, k \in \mathbb{Z}$ and $l \in \mathbb{Z}^*$ subject to the following relations

$$\begin{aligned} K_i K_i^{-1} &= K_i^{-1} K_i = 1, & [K_i, K_j] &= [K_i, h_{j,l}] = [h_{i,l}, h_{j,k}] = 0, \\ K_i x_{j,k}^\pm K_i^{-1} &= v^{\pm a_{ij}} x_{j,k}^\pm, \\ [h_{i,l}, x_{j,k}^\pm] &= \pm \frac{1}{l} [a_{ij}] x_{j,k+l}^\pm, \\ x_{i,k+1}^\pm x_{j,l}^\pm - v^{\pm a_{ij}} x_{j,l}^\pm x_{i,k+1}^\pm &= v^{\pm a_{ij}} x_{i,k}^\pm x_{j,l+1}^\pm - x_{j,l+1}^\pm x_{i,k}^\pm, \\ [x_{i,k}^+, x_{j,l}^-] &= \delta_{ij} \frac{\psi_{i,k+l} - \varphi_{i,k+l}}{v - v^{-1}}, \end{aligned}$$

For $i \neq j$ and $n = 1 - a_{ij}$,

$$\text{Sym}_{k_1, \dots, k_n} \sum_{t=0}^{1-a_{ij}} (-1)^t \begin{bmatrix} n \\ t \end{bmatrix} x_{i,k_1}^\pm \cdots x_{i,k_t}^\pm x_{j,l}^\pm x_{i,k_{t+1}}^\pm \cdots x_{i,k_n}^\pm = 0$$

where $\text{Sym}_{k_1, \dots, k_n}$ denotes symmetrization with respect to the indices $k_1, \dots, k_n$, and where $\psi_{i,k}$ and $\varphi_{i,k}$ are defined by the following equations :

$$\begin{aligned} \sum_{k \geq 0} \psi_{i,k} u^k &= K_i \exp((v - v^{-1}) \sum_{k=1}^{\infty} h_{i,k} u^k), \\ \sum_{k \geq 0} \varphi_{i,k} u^k &= K_i^{-1} \exp(-(v - v^{-1}) \sum_{k=1}^{\infty} h_{i,-k} u^{-k}). \end{aligned}$$

1.9. The quantum loop algebra $\mathbf{U}_v(L\mathfrak{gl}_n)$ is the $\mathbb{C}(v)$ -algebra generated by $U_v(L\mathfrak{sl}_n)$ and generators $K_0, K_0^{-1}, h_{0,l}$ for $l \in \mathbb{Z}^*$ with relations

$$\begin{aligned} K_0 K_0^{-1} &= K_0^{-1} K_0 = 1, \\ [K_0, K_i] &= [K_0, h_{i,l}] = [K_0, h_{0,l}] = [h_{0,l}, h_{i,k}] = [h_{0,l}, h_{0,k}] = 0, \\ K_0 x_{i,k}^\pm K_0^{-1} &= v^{\mp \delta_{i1}} x_{i,k}^\pm, \\ [h_{0,l}, x_{i,k}^\pm] &= \mp \delta_{i,1} \frac{[l]}{l} x_{i,l+k}^\pm. \end{aligned}$$

2. WEIGHTED PROJECTIVE LINES

We recall in this section the definition and main properties of the weighted projective lines $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ and the categories of coherent sheaves on them, as studied in [GL].

2.1. Let k be an arbitrary field and let $\mathbf{p} = \{p_1, \dots, p_n\} \in (\mathbb{N}^*)^n$. Consider the $\mathbb{Z}$ -module

$$L(\mathbf{p}) = \mathbb{Z}\vec{x}_1 \oplus \dots \oplus \mathbb{Z}\vec{x}_n / J$$

where J is the $\mathbb{Z}$ -submodule generated by $\{p_1 \vec{x}_1 - p_s \vec{x}_s \mid s = 2, \dots, n\}$. Set $\vec{c} = p_1 \vec{x}_1 = \dots = p_n \vec{x}_n$. Then $L(\mathbf{p})/\mathbb{Z}\vec{c} \simeq \prod_{s=1}^n \mathbb{Z}/p_s \mathbb{Z}$ and we can consider $L(\mathbf{p})$ as a rank one abelian group. We set $L^+(\mathbf{p}) = \{l_1 \vec{x}_1 + \dots + l_n \vec{x}_n \mid l_i \geq 0\}$ and we write $\vec{x} \geq \vec{y}$ if $\vec{x} - \vec{y} \in L^+(\mathbf{p})$.

Let $k[L(\mathbf{p})]$ be the group Hopf algebra of $L(\mathbf{p})$ and let

$$G(\mathbf{p}) = \text{Spec } k[L(\mathbf{p})] \subset (k^*)^n$$

be the associated affine algebraic group. Hence the $\bar{k}$ -points of $G(\mathbf{p})$ are

$$G(\mathbf{p})(\bar{k}) = \{(t_1, \dots, t_n) \in (\bar{k}^*)^n \mid t_1^{p_1} = \dots = t_n^{p_n}\}.$$

By definition $G(\mathbf{p})$ acts on k^n and induces on $S(\mathbf{p}) := k[X_1, \dots, X_n]$ the structure of an $L(\mathbf{p})$ -graded algebra such that $\deg X_s = \vec{x}_s$.

Let $\underline{\lambda} = \{\lambda_1, \dots, \lambda_n\}$ be a collection of distinct closed points (of degree 1) of $\mathbb{P}^1(k)$, normalized in such a way that $\lambda_1 = \infty$, $\lambda_2 = 0$ and $\lambda_3 = 1$. Let $I(\mathbf{p}, \underline{\lambda})$ be the $L(\mathbf{p})$ -graded ideal in $S(\mathbf{p})$ generated by the polynomials $X_s^{p_s} - (X_2^{p_2} - \lambda_s X_1^{p_1})$ for $s = 3, \dots, n$ and set $S(\mathbf{p}, \underline{\lambda}) = S(\mathbf{p})/I(\mathbf{p}, \underline{\lambda})$. We will denote by $x_1, \dots, x_n$ the images of $X_1, \dots, X_n$ in $S(\mathbf{p}, \underline{\lambda})$. It is known that $S(\mathbf{p}, \underline{\lambda})$ is a graded factorial domain of Krull dimension equal to 2. We let

$$\mathbb{X}_{\mathbf{p}, \underline{\lambda}} = \text{Specgr } S(\mathbf{p}, \underline{\lambda})$$

be the set of prime homogeneous ideals in $S(\mathbf{p}, \underline{\lambda})$.

Proposition 2.1 ([GL]). *The nonzero homogeneous prime elements in $S(\mathbf{p}, \underline{\lambda})$ are of the form*

- i) $x_1, \dots, x_n$ (the exceptional points), or
- ii) $F(x_1^{p_1}, x_2^{p_2})$ where $F \in k[T_1, T_2]$ is a prime homogeneous polynomial (ordinary points).

We will denote by σ_s , $s = 1, \dots, n$ the closed points of $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ corresponding to the primes $x_1, \dots, x_n$. Let $p = \text{l.c.m.}(p_1, \dots, p_n)$. We define the *degree* of a closed point x by $\deg(\sigma_i) = \frac{p}{p_i}$ for exceptional points and $\deg(x) = pd$ if x is associated to a prime homogeneous polynomial $P(x_1^{p_1}, x_2^{p_2})$ of degree d .

For any homogeneous $f \in S(\mathbf{p}, \underline{\lambda})$ we set $V_f = \{\mathfrak{p} \in \mathbb{X}_{\mathbf{p}, \underline{\lambda}} \mid f \in \mathfrak{p}\}$ and $D_f = \mathbb{X}_{\mathbf{p}, \underline{\lambda}} \setminus V_f$. The sets D_f form a basis of the Zariski topology on $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$.

2.2. Let $\mathcal{O}_{\mathbb{X}}$ be the sheaf of $L(\mathbf{p})$ -graded algebras on $\mathbb{X}_{\mathbf{p},\underline{\lambda}}$ associated to the presheaf $D_f \mapsto S(\mathbf{p},\underline{\lambda})_f$ where

$$S(\mathbf{p},\underline{\lambda})_f = \left\{ \frac{g}{f^l} \mid g \in S(\mathbf{p},\underline{\lambda}), l \in \mathbb{N} \right\}$$

is the localization of $S(\mathbf{p},\underline{\lambda})$ at f . Let us denote by $\mathcal{O}_{\mathbb{X}}\text{-Mod}$ the category of sheaves of $L(\mathbf{p})$ -graded $\mathcal{O}_{\mathbb{X}}$ -modules on $\mathbb{X}_{\mathbf{p},\underline{\lambda}}$.

For any $\vec{x} \in L(\mathbf{p})$ and any object $\mathcal{M} = \bigoplus_{\vec{l} \in L(\mathbf{p})} \mathcal{M}[\vec{l}]$ in $\mathcal{O}_{\mathbb{X}}\text{-Mod}$ we denote by $\mathcal{M}(\vec{x})$ the shift of $\mathcal{M}$ by $\vec{x}$ (i.e $\mathcal{M}(\vec{x})[\vec{l}] := \mathcal{M}[\vec{x} + \vec{l}]$). The category $\text{Coh}(\mathbb{X}_{\mathbf{p},\underline{\lambda}})$ is now defined as the full subcategory of $\mathcal{O}_{\mathbb{X}}\text{-Mod}$ consisting of sheaves $\mathcal{M}$ for which there exists an open covering $\{U_i\}$ of $\mathbb{X}_{\mathbf{p},\underline{\lambda}}$ and, for each i , an exact sequence

$$\bigoplus_{s=1}^N \mathcal{O}_{\mathbb{X}}(\vec{l}_s)|_{U_i} \rightarrow \bigoplus_{t=1}^M \mathcal{O}_{\mathbb{X}}(\vec{k}_t)|_{U_i} \rightarrow \mathcal{M}|_{U_i} \rightarrow 0.$$

Note that for any $\mathcal{M} \in \text{Coh}(\mathbb{X}_{\mathbf{p},\underline{\lambda}})$ the space $\Gamma(\mathbb{X}_{\mathbf{p},\underline{\lambda}}, \mathcal{M})$ of global sections of $\mathcal{M}$ is an $L(\mathbf{p})$ -graded $S(\mathbf{p},\underline{\lambda})$ -module.

Proposition 2.2 ([GL]). *The category $\text{Coh}(\mathbb{X}_{\mathbf{p},\underline{\lambda}})$ is hereditary, i.e*

$$\text{Ext}^2(\mathcal{M}, \mathcal{N}) = 0$$

for any $\mathcal{M}, \mathcal{N}$. Moreover, for any $\mathcal{M}, \mathcal{N}$, $\text{Hom}(\mathcal{M}, \mathcal{N})$ and $\text{Ext}^1(\mathcal{M}, \mathcal{N})$ are finite-dimensional.

We now define the *support* of a sheaf. Let x be any closed point of $\mathbb{X}_{\mathbf{p},\underline{\lambda}}$. Let us denote by $\mathfrak{m}_x$ the corresponding maximal ideal and set

$$\mathcal{O}_{\mathbb{X},x} = \left\{ \frac{f}{g} \mid f, g \in S(\mathbf{p},\underline{\lambda}), g \text{ homogeneous}, g \notin \mathfrak{m}_x \right\}.$$

This is an $L(\mathbf{p})$ -graded discrete valuation ring. If $\mathcal{M} \in \text{Coh}(\mathbb{X}_{\mathbf{p},\underline{\lambda}})$ we define $\mathcal{M}_x = \mathcal{M}(D_f) \otimes_{\mathcal{O}_{\mathbb{X}}(D_f)} \mathcal{O}_{\mathbb{X},x}$ where D_f is any neighborhood of x . The support of $\mathcal{M}$ is the Zariski closure of the set $\{x \in \mathbb{X}_{\mathbf{p},\underline{\lambda}} \mid \mathcal{M}_x \neq 0\}$.

Let $\xi \in \mathbb{X}_{\mathbf{p},\underline{\lambda}}$ denote the generic point, associated to the prime ideal (0). Set

$$\mathcal{O}_{\mathbb{X},\xi} = \left\{ \frac{f}{g} \mid f, g \in S(\mathbf{p},\underline{\lambda}), g \text{ homogeneous}, g \neq 0 \right\}.$$

This $L(\mathbf{p})$ -graded ring is Morita equivalent to its degree zero component $\mathcal{O}_{\mathbb{X},\xi}[0] \simeq k\left(\frac{x_1^{p_1}}{x_2^{p_2}}\right)$. The *rank* of a coherent sheaf $\mathcal{M}$ is defined as

$$\mathbf{r}(\mathcal{M}) = \dim \mathcal{M}(D_f) \otimes_{\mathcal{O}_{\mathbb{X}}(D_f)} \mathcal{O}_{\mathbb{X},\xi}$$

for any D_f .

2.3. Let $S(\mathbf{p},\underline{\lambda})\text{-modgr}$ be the category of finitely generated $L(\mathbf{p})$ -graded $S(\mathbf{p},\underline{\lambda})$ -modules. As in the classical case, one can introduce a sheafification functor

$$\begin{aligned} Sh : S(\mathbf{p},\underline{\lambda})\text{-modgr} &\rightarrow \text{Coh}(\mathbb{X}_{\mathbf{p},\underline{\lambda}}) \\ M &\mapsto \tilde{M} \end{aligned}$$

where $\tilde{M}$ is the sheaf associated to the presheaf

$$D_f \mapsto M_f = \left\{ \frac{m}{f^l} \mid m \in M, l \in \mathbb{N} \right\}.$$

Let $L^+(\mathbf{p}) \subset L(\mathbf{p})$ be the monoid generated by $\vec{x}_1, \dots, \vec{x}_n$. Denote by $S^+(\mathbf{p},\underline{\lambda})\text{-modgr}$ be the full subcategory of $S(\mathbf{p},\underline{\lambda})\text{-modgr}$ consisting of $L^+(\mathbf{p})$ -graded modules.

Consider the global sections functor

$$\Gamma_+ : \text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}}) \rightarrow S^+(\mathbf{p}, \underline{\lambda}) - \text{modgr}$$

$$\mathcal{M} \mapsto \bigoplus_{l \geq 0} \Gamma(X, \mathcal{M})[\vec{l}].$$

The following analogue of Serre's theorem holds :

Proposition 2.3 ([GL]). *The functor Γ_+ defines a full embedding and for any $\mathcal{M} \in \text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$ we have $\text{Sh}(\Gamma_+(\mathcal{M})) \simeq \mathcal{M}$. The functor Sh is exact and induces an equivalence*

$$\text{Sh} : S(\mathbf{p}, \underline{\lambda}) - \text{modgr} / S(\mathbf{p}, \underline{\lambda}) - \text{modgr}_0 \xrightarrow{\sim} \text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}}),$$

where $S(\mathbf{p}, \underline{\lambda}) - \text{modgr}_0$ denotes the quotient of $S(\mathbf{p}, \underline{\lambda}) - \text{modgr}$ by the Serre subcategory of finite-length modules.

In particular, we have

$$(2.1) \quad \text{Hom}(\mathcal{O}_{\mathbb{X}}(\vec{x}), \mathcal{O}_{\mathbb{X}}(\vec{y})) = S(\mathbf{p}, \underline{\lambda})[\vec{y} - \vec{x}].$$

2.4. A sheaf $\mathcal{M}$ is *locally free* or is a *vector bundle* if there exists an open covering $\{U_i\}$ of $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ and, for each i , an isomorphism $\bigoplus_{i=1}^N \mathcal{O}_{\mathbb{X}}(\vec{l}_i)|_{U_i} \xrightarrow{\sim} \mathcal{M}|_{U_i}$ for some suitable $\vec{l}_i$. Every rank one locally free sheaf on $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ is isomorphic to $\mathcal{O}_{\mathbb{X}}(\vec{x})$ for some $\vec{x} \in L(\mathbf{p})$. The full subcategory consisting of locally free sheaves is stable under extensions (but not under quotient, and is not abelian).

A sheaf $\mathcal{M}$ is a *torsion sheaf* if it is a finite length object in $\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$, or, equivalently, if it is of rank zero. By definition, the full subcategory $\mathcal{T}$ consisting of torsion sheaves is a Serre subcategory. The following properties can be found in [GL] :

- i) Every sheaf $\mathcal{M} \in \text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$ can be decomposed as $\mathcal{M} = \mathcal{M}_t \oplus \mathcal{M}'$ for a (unique) $\mathcal{M}_t \in \mathcal{T}$ and some $\mathcal{M}' \in \mathcal{F}$.
- ii) $\text{Hom}(\mathcal{M}_t, \mathcal{M}_f) = \text{Ext}^1(\mathcal{M}_f, \mathcal{M}_t) = 0$ for every $\mathcal{M}_t \in \mathcal{T}$ and $\mathcal{M}_f \in \mathcal{F}$.

The category $\mathcal{T}$ decomposes as a product $\mathcal{T} = \sqcup_{x \in \mathbb{X}_{\mathbf{p}, \underline{\lambda}}} \mathcal{T}_x$ where $\mathcal{T}_x$ is the category of torsion sheaves with support at the closed point $x \in \mathbb{X}_{\mathbf{p}, \underline{\lambda}}$. Moreover, $\mathcal{T}_x$ is Morita-equivalent to the category $\text{Mod}_0 \mathcal{O}_{\mathbb{X}, x}$ of finite-length $L(\mathbf{p})$ -graded $\mathcal{O}_{\mathbb{X}, x}$ -modules.

These can be explicitly described. Let $\mathcal{C}_l$ denote quiver of type $A_{l-1}^{(1)}$ with cyclic orientation. We let $\mathcal{C}_1$ be the quiver with one vertex and one arrow. A representation of $\mathcal{C}_p$ over a field k is by definition a collection of k -vector spaces V_i with $i \in \mathbb{Z}/p\mathbb{Z}$ together with a collection of linear maps $x_i : V_i \rightarrow V_{i-1}$. Morphisms between two representations (V_i, x_i) and (W_i, y_i) are k -linear maps $\phi_i : V_i \rightarrow W_i$ such that $\phi_i x_i = y_i \phi_i$. Finally, a representation (V_i, x_i) is called nilpotent if there exists $N \gg 0$ such that $x_{i+N} \cdots x_i = 0$ for all i .

Lemma 2.1. *The following holds :*

- i) *Let x be an ordinary closed point of degree d (see Proposition 2.1) and let k_x denote the residue field at x . Then $\mathcal{T}_x$ is Morita-equivalent to the category of nilpotent representation of the cyclic quiver $\mathcal{C}_1$ over k_x .*
- ii) *Let $1 \leq s \leq n$. The category $\mathcal{T}_{\sigma_s}$ is Morita equivalent to the category of nilpotent representations of $\mathcal{C}_{p^s}$ over k .*

In case i) of the lemma above, the simple module S_x can be realized in the following way : let $\pi_x \in S(\mathbf{p}, \Delta)$ denote the prime corresponding to x . Multiplication by π_x leads to an exact sequence

$$0 \rightarrow \mathcal{O}_{\mathbb{X}} \rightarrow \mathcal{O}_{\mathbb{X}}(d\vec{c}) \rightarrow S_x \rightarrow 0.$$

Moreover, for any $\vec{k} \in L(\mathbf{p})$ we have $S_x(\vec{k}) \simeq S_x$. In case ii), the simple sheaves $S_j^{(s)}$, $j = 1, \dots, p_s$ with support at σ_s can be realized as follows : multiplication by x_s leads to exact sequences

$$0 \rightarrow \mathcal{O}_{\mathbb{X}}((j-1)\vec{x}_s) \rightarrow \mathcal{O}_{\mathbb{X}}(j\vec{x}_s) \rightarrow S_j^{(s)} \rightarrow 0.$$

Moreover, for any $\vec{k} = \sum_i k_i \vec{x}_i \in L(\mathbf{p})$ we have

$$(2.2) \quad S_j^{(s)}(\vec{k}) \simeq S_{j+k_s \bmod p_s}^{(s)}.$$

In particular, $S_j^{(s)}(t\vec{c}) \simeq S_j^{(s)}$ for any $t \in \mathbb{Z}$. Finally, we note the following result :

Proposition 2.4. *Let $K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta}))$ be the Grothendieck group of $\text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta})$ and let $[\mathcal{M}]$ denote the class of a coherent sheaf $\mathcal{M}$. Then*

$$K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta})) \simeq (\mathbb{Z}[\mathcal{O}_{\mathbb{X}}] \oplus \mathbb{Z}[\mathcal{O}_{\mathbb{X}}(\vec{c})] \bigoplus_{s,j} \mathbb{Z}[S_j^{(s)}]) / I$$

where I is the subgroup generated by $\{\sum_j [S_j^{(s)}] + [\mathcal{O}_{\mathbb{X}}] - [\mathcal{O}_{\mathbb{X}}(\vec{c})]\}_{s=1, \dots, n}$.

2.5. Let $\vec{\omega} = (n-1)\vec{c} - \sum_i \vec{x}_i$.

Theorem 2.1 (Serre duality, [GL]). *Let $\mathcal{M}, \mathcal{N} \in \text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta})$. There is a functorial isomorphism*

$$(\text{Ext}^1(\mathcal{M}, \mathcal{N}))^* \xrightarrow{\sim} \text{Hom}(\mathcal{N}, \mathcal{M}(\vec{\omega})).$$

2.6. Define a group homomorphism $\partial : L(\mathbf{p}) \rightarrow \mathbb{Z}$ by setting $\partial(\vec{x}_s) = \frac{p}{p_s}$. There is a unique *degree map* $\mathbf{d} : K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta})) \rightarrow \mathbb{Z}$ satisfying $\mathbf{d}([\mathcal{O}_{\mathbb{X}}(\vec{x})]) = \partial(\vec{x})$ and $\mathbf{d}([S]) = \deg(x)$ if S is a simple torsion sheaf supported at a point x .

Now for any $\mathcal{M} \in \text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta})$ set $\chi(\mathcal{M}) = \dim \text{Hom}(\mathcal{O}_{\mathbb{X}}, \mathcal{M}) - \dim \text{Ext}^1(\mathcal{O}_{\mathbb{X}}, \mathcal{M})$ (the *Euler characteristic* of $\mathcal{M}$), and put

$$\overline{\chi}(\mathcal{M}) = \sum_{k=0}^{p-1} \chi(\mathcal{M}(-k\vec{\omega})).$$

Theorem 2.2 (Riemann-Roch theorem, [GL]). *For any $\mathcal{M} \in \text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta})$ we have*

$$\overline{\chi}(\mathcal{M}) = \mathbf{r}(\mathcal{M})\overline{\chi}(\mathcal{O}_{\mathbb{X}}) + \mathbf{d}(\mathcal{M}),$$

and $\overline{\chi}(\mathcal{O}_{\mathbb{X}}) = -\frac{p}{2}\partial(\vec{\omega})$.

The previous theorem motivates the following definition of the genus of $\mathbb{X}_{\mathbf{p}, \Delta}$:

$$g(\mathbb{X}_{\mathbf{p}, \Delta}) = 1 + \frac{1}{2}\partial(\vec{\omega}).$$

Using this notation, the Riemann-Roch theorem admits the following corollary. We set $\langle M, N \rangle = \dim \text{Hom}(M, N) - \dim \text{Ext}^1(M, N)$. Then, for all coherent sheaves M and N ,

$$(2.3) \quad \sum_{k=0}^{p-1} \langle M(k\vec{\omega}), N \rangle = p(1 - g_{\mathbb{X}_{\mathbf{p}, \Delta}})\mathbf{r}(M)\mathbf{r}(N) + \mathbf{r}(M)\mathbf{d}(N) - \mathbf{r}(N)\mathbf{d}(M).$$

Let us associate to the sequence $(p_1, \dots, p_l)$ the “star Dynkin diagram” $\mathbb{T}_{p_1, \dots, p_n}$ (see Section 1.3). The genus of $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ and the properties of $\mathbb{T}_{p_1, \dots, p_n}$ are related in the following way :

- i) $g(\mathbb{X}_{\mathbf{p}, \underline{\lambda}}) < 1$ if and only if $\mathbb{T}_{p_1, \dots, p_n}$ is a finite Dynkin diagram,
- ii) $g(\mathbb{X}_{\mathbf{p}, \underline{\lambda}}) = 1$ if and only if $\mathbb{T}_{p_1, \dots, p_n}$ is an affine Dynkin diagram.

2.7. One of our main tools in the study of the category $\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$ will be the Harder-Narasimhan (HN) filtration, introduced in [GL]. Define the slope of a nonzero coherent sheaf $\mathcal{F} \in \text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$ as $\mu(\mathcal{F}) = \frac{\mathbf{d}(\mathcal{F})}{\mathbf{r}(\mathcal{F})} \in \mathbb{Q} \cup \{\infty\}$ (where $\mathbf{r}(\mathcal{F})$ and $\mathbf{d}(\mathcal{F})$ are the rank and degree of $\mathcal{F}$). Note that $\mu(\mathcal{F}) = \infty$ if and only if $\mathcal{F}$ is a torsion sheaf. The sheaf $\mathcal{F}$ is called stable (resp. semistable) if for every subsheaf $\mathcal{G} \subset \mathcal{F}$ we have $\mu(\mathcal{G}) < \mu(\mathcal{F})$ (resp. $\mu(\mathcal{G}) \leq \mu(\mathcal{F})$). For any $q \in \mathbb{Q}$ let $\mathcal{C}_q$ be the full subcategory of $\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$ consisting the zero sheaf together with all sheaves of slope q .

Proposition 2.5 ([GL]). *For each $q \in \mathbb{Q}$ the subcategory $\mathcal{C}_q$ is abelian and closed under extensions. Moreover, each object is finite length and the simple objects are the stable bundles. Finally, $\text{Hom}(\mathcal{F}, \mathcal{G}) = 0$ if $\mathcal{F} \in \mathcal{C}_q$, $\mathcal{G} \in \mathcal{C}_{q'}$ and $q > q'$.*

The relevance of this notion for us is then explained by the following results.

Proposition 2.6 ([GL]). *If Γ is finite (resp. affine) then every indecomposable locally free sheaf is stable (resp. semistable).*

From Serre duality and the fact that $\partial(\vec{\omega}) \leq 0$ if $g(\mathbb{X}_{\mathbf{p}, \underline{\lambda}}) \leq 1$ and $\partial(\vec{\omega}) = 0$ if $g(\mathbb{X}_{\mathbf{p}, \underline{\lambda}}) = 1$ we deduce

Corollary 2.1. *Suppose that Γ is finite. Then*

$$q \leq q' \Rightarrow \text{Ext}^1(\mathcal{F}, \mathcal{G}) = 0 \text{ for all } \mathcal{F} \in \mathcal{C}_q, \mathcal{G} \in \mathcal{C}_{q'}.$$

Suppose that Γ is affine. Then

$$q < q' \Rightarrow \text{Ext}^1(\mathcal{F}, \mathcal{G}) = 0 \text{ for all } \mathcal{F} \in \mathcal{C}_q, \mathcal{G} \in \mathcal{C}_{q'}.$$

3. RINGEL-HALL ALGEBRAS

3.1. Let k be a finite field with q elements and let $\mathcal{A}$ be a (small) abelian k -linear category. We assume the following :

- i) $\mathcal{A}$ is hereditary, i.e $\text{Ext}^2(V, W) = 0$ for any objects V and W ,
- ii) $\mathcal{A}$ is Hom and Ext-finite, i.e for any objects V, W the spaces $\text{Hom}(V, W)$ and $\text{Ext}^1(V, W)$ are finite-dimensional.

Let $\text{Iso}(\mathcal{A})$ denote the set of isomorphism classes of objects in $\mathcal{A}$ and let $K_0(\mathcal{A})$ be the Grothendieck group of $\mathcal{A}$. The assignment

$$\begin{aligned} \text{Iso}(\mathcal{A}) \times \text{Iso}(\mathcal{A}) &\rightarrow \mathbb{Z} \\ \langle V, W \rangle &\mapsto \dim \text{Hom}(V, W) - \dim \text{Ext}^1(V, W) \end{aligned}$$

descends to a well-defined biadditive form $\langle \cdot, \cdot \rangle : K_0(\mathcal{A}) \otimes_{\mathbb{Z}} K_0(\mathcal{A}) \rightarrow \mathbb{Z}$ called the *Euler form*.

3.2. Let $S, T, U \in \text{Iso}(\mathcal{A})$. Let $\mathcal{P}_{S,T}^U$ be the (finite) set of all exact sequences

$$0 \rightarrow T \xrightarrow{f} U \xrightarrow{g} S \rightarrow 0.$$

The group $\text{Aut}(S) \times \text{Aut} T$ acts faithfully on $\mathcal{P}_{S,T}^U$ and we set

$$(3.1) \quad \mathbf{P}_{S,T}^U = |\mathcal{P}_{S,T}^U / \text{Aut}(S) \times \text{Aut}(T)| \in \mathbb{N}.$$

Consider the ring $\mathbb{C}_q := \mathbb{C}[v, v^{-1}]/(v^2 - q)$. We endow the free $\mathbb{C}_q$ -module

$$\mathbf{H}_{\mathcal{A}} := \bigoplus_{S \in \text{Iso}(\mathcal{A})} \mathbb{C}_q[S]$$

with an associative algebra structure by setting

$$(3.2) \quad [S] \cdot [T] := v^{\langle S, T \rangle} \sum_U \mathbf{P}_{S,T}^U [U].$$

Such algebras were first considered by Hall ([Hal], see Section 3.3) when $\mathcal{A}$ is the category of representations of a discrete valuation ring with finite residue field, and later in much more generality by Ringel (see e.g [Ri1]).

4. RINGEL-HALL ALGEBRAS OF CYCLIC QUIVERS

4.1. Let us consider the case of the quiver $\mathcal{C}_1$. The isomorphism classes of indecomposable (nilpotent) representations of $\mathcal{C}_1$ over the finite field $\mathbb{F}_q$ with q elements is in natural bijection with $\mathbb{N}^*$ and we denote by $|n\rangle$ the class of the indecomposable representation of dimension n . The isomorphism classes of (nilpotent) representations is thus in bijection with the set Π of all partitions via the assignment $\lambda = (\lambda_1, \dots, \lambda_r) \mapsto |\lambda\rangle = |\lambda_1\rangle \oplus \dots \oplus |\lambda_r\rangle$. Let $\mathbf{H}_1^{(q)}$ be the Hall algebra of $\mathcal{C}_1$ over $\mathbb{F}_q$ and set $\mathbf{e}_r = v^{r(r-1)} |(1^r)\rangle$. It is known that the structure constants for $\mathbf{H}_1^{(q)}$ are polynomials in q and one can consider $\mathbf{H}_1^{(q)}$ as a $\mathbb{C}_v$ -algebra.

Let $\Gamma = \mathbb{C}[y_1, y_2, \dots]^{\oplus \infty}$ be Macdonald's ring of symmetric functions (see [M]) and let $\{e_r\}_{r \in \mathbb{N}}$ denote the elementary symmetric polynomial. The following well-known result is due to P. Hall.

Theorem 4.1. *The assignment $e_r \mapsto \mathbf{e}_r$ uniquely extends to an isomorphism of algebras $\Upsilon : \Gamma \otimes_{\mathbb{C}} \mathbb{C}_v \xrightarrow{\sim} \mathbf{H}_1^{(q)}$.*

Under this isomorphism the element $|\lambda\rangle$ corresponds to $q^{-n(\lambda)} P_{\lambda}(q^{-1})$ where $n(\lambda) = \sum_i (i-1)\lambda_i$ and $P_{\lambda}(t)$ is the Hall-Littlewood polynomial (see [M]). Let $p_r \in \Gamma$ denote the power-sum symmetric function and set

$$\mathbf{h}_r = \frac{[r]}{r} \Upsilon(p_r).$$

It follows from [M], III.7, Ex. 2, that

$$(4.1) \quad \mathbf{h}_r = \frac{[r]}{r} \sum_{\lambda, |\lambda|=r} n(l(\lambda) - 1) |\lambda\rangle,$$

where $n(l) = \prod_{i=1}^l (1 - v^{2i})$.

4.2. Let $p \in \mathbb{N}$ and let $\mathbf{H}_p^{(q)}$ be the Hall algebra of the cyclic quiver $\mathcal{C}_p$ over $\mathbb{F}_q$ (we will also denote it simply by $\mathbf{H}_p$ when there is no ambiguity in the ground field). Denote by $\{\epsilon_i\}_{i \in \mathbb{Z}/p\mathbb{Z}}$ the canonical basis of $\mathbb{Z}^{\mathbb{Z}/p\mathbb{Z}}$. For each $i \in \mathbb{Z}/p\mathbb{Z}$ and $l \in \mathbb{N}$ define the *cyclic segment* $[i; l]$ to be the image of the projection to $\mathbb{Z}/p\mathbb{Z}$ of the segment $[i' - (l - 1), i']$ for any $i' \in \mathbb{Z}$, $i' \equiv i \pmod{p}$. A *cyclic multisegment* is a finite linear combination $\mathbf{m} = \sum_{i,l} a_{il} [i; l]$ with $a_{il} \in \mathbb{N}$. The isomorphism classes of representations (resp. indecomposable representations) of $\mathcal{C}_p$ over k are in natural bijection with the set of cyclic multisegments (resp. cyclic segments). For $i \in \mathbb{Z}/p\mathbb{Z}$ we let $S_i = [[i; 1]]$ be the class of the corresponding simple object. More generally, for any $n \geq 1$ we denote by $S_i^{(n)}$ the class of the semisimple module $[i; 1]^{\oplus n}$. Finally, let $d(\mathbf{m}) = \sum_{i,l} a_{il} (\epsilon_i + \dots + \epsilon_{i-(l-1)}) \in \mathbb{N}^{\mathbb{Z}/p\mathbb{Z}}$ be the *degree* of a multisegment $\mathbf{m}$ and set $\delta = (1, \dots, 1)$. The algebra $\mathbf{H}_p$ is graded and we denote by $\mathbf{H}_p[d]$ the piece of degree d . It follows from [Ril] that the assignment $E_i^{(n)} \mapsto v^{n(n-1)/2} [S_i^{(n)}]$ defines an embedding $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \hookrightarrow \mathbf{H}_p$.

Theorem 4.2 ([S]). *The embedding $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \hookrightarrow \mathbf{H}_p$ extends to an isomorphism*

$$\mathbf{H}_p \simeq \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \otimes_{\mathbb{C}_v} \mathcal{Z}$$

where $\mathcal{Z} = \mathbb{C}_v[x_1, x_2, \dots]$ is a central subalgebra and where the element x_i is homogeneous of degree $i\delta$.

Remark. The algebra structure $\bullet : \mathbf{H}_p \otimes \mathbf{H}_p \rightarrow \mathbf{H}_p$ used in [S] differs slightly from the one defined by (3.2). Namely, if $f \in \mathbf{H}_p[d]$ and $f' \in \mathbf{H}_p[d']$ then

$$f \bullet f' = v^{-2 \sum_i d_i d'_i} f' \cdot f.$$

Note also that the opposite orientation of the cyclic quiver is used in [S].

To any $(\lambda_1, \dots, \lambda_r) \in \Pi$ we associate

$$f_\lambda = \sum_j [0; p\lambda_j] \in \mathbf{H}_p$$

and we set

$$(4.2) \quad \mathbf{H}_p^0 := \bigoplus_{\lambda \in \Pi} \mathbb{C}_v f_\lambda$$

The assignment $|\lambda\rangle \mapsto f_\lambda$ uniquely extends to an algebra isomorphism $\Psi : \mathbf{H}_1^{(q)} \xrightarrow{\sim} \mathbf{H}_p^0$. In particular, $\mathbf{H}_p^0$ is a commutative subalgebra of $\mathbf{H}_p$ and is freely generated by any of the three sets $\{\mathbf{e}_r\}_{r \in \mathbb{N}^*}$, $\{\mathbf{l}_r\}_{r \in \mathbb{N}^*}$ and $\{\mathbf{h}_r\}_{r \in \mathbb{N}^*}$ where

$$\mathbf{e}_r = \Psi(\mathbf{e}_r), \quad \mathbf{l}_r = f_{(r)}, \quad \mathbf{h}_r = \Psi(\mathbf{h}_r).$$

Lemma 4.1. *The multiplication maps $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \otimes \mathbf{H}_p^0 \rightarrow \mathbf{H}_p$ and $\mathbf{H}_p^0 \otimes \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \rightarrow \mathbf{H}_p$ induce isomorphisms of $\mathbb{C}_v$ -modules.*

Proof. It is enough to prove both statements for the algebra structure $\bullet$ used in [S]. We first recall some notation and results from [S]. By [S], Proposition 2.2, there exists a canonical isomorphism $i : \Gamma \otimes \mathbb{C}_v \xrightarrow{\sim} \mathcal{Z}$. Let $\{s_\lambda\}_{\lambda \in \Pi}$ denote the Schur basis of Γ and set $\mathbf{s}_\lambda = i(s_\lambda)$. Let $\mathbf{b}_\mathbf{m}$ be the canonical basis of $\mathbf{H}_p$. To a partition $\lambda \in \Pi$ we associate the multisegments $\mathbf{m}(\lambda) = \sum_i [1 - i, \lambda_i - i]$ and $\mathbf{m}^\lambda = \sum_{i,j} [i, \lambda_j]$. We have the following (see [S], Proposition 2.4)

$$(4.3) \quad \sum_i E_i \bullet \mathbf{H}_p = \bigoplus_{\mathbf{m} \notin \{\mathbf{m}^\lambda, \lambda \in \Pi\}} \mathbb{C}_v \mathbf{b}_\mathbf{m}.$$

Let Λ^∞ be the level one Fock space of $\mathbf{H}_p$ considered in [S] and let $|0\rangle$ be its vacuum vector. Let $\{\mathbf{b}_\lambda^\pm \mid \lambda \in \Pi\}$ be the Leclerc-Thibon canonical bases of Λ^∞ (see [LT]). Then (see [S], Proposition 3.1 and [LT], Theorem 6.9)

$$(4.4) \quad i)\mathbf{b}_{\mathbf{m}(\lambda)}|0\rangle = \mathbf{b}_\lambda^+, \quad ii)x \cdot |0\rangle = 0 \Leftrightarrow x \in \bigoplus_{\mathbf{m} \notin \{\mathbf{m}(\lambda), \lambda \in \Pi\}} \mathbb{C}_v \mathbf{b}_{\mathbf{m}}$$

$$(4.5) \quad \mathbf{s}_\lambda|0\rangle = \mathbf{b}_{p\lambda}^-.$$

Finally, it is easy to prove that

$$(4.6) \quad \mathbb{C}_v[\mathbf{l}_1, \mathbf{l}_2, \dots]|0\rangle = \bigoplus_{\mu} \mathbb{C}_v|p\mu\rangle.$$

We now prove the lemma by induction. A direct computation shows that $\mathbf{H}_p[\delta] = \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p)[\delta] \oplus \mathbb{C}_v \mathbf{l}_1$. Assume that for any $r < k$ we have

$$\mathbf{H}_p[r\delta] = (\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \otimes \mathbb{C}_v[\mathbf{l}_1, \dots, \mathbf{l}_r])[r\delta].$$

Observe that $\mathbf{b}^\lambda|0\rangle = \mathbf{b}_{n\lambda}^+$ and recall that for any μ , $\mathbf{b}_\mu^\pm \in |\mu\rangle + \bigoplus_{\nu < \mu} \mathbb{C}_v|\nu\rangle$. From (4.4)i), (4.5) and (4.6) it now follows that there exists $P_k \in \mathbb{C}_v[\mathbf{l}_1, \dots, \mathbf{l}_k]$ and $x \in \bigoplus_{\mathbf{m} \neq \mathbf{m}^\lambda} \mathbb{C}_v \mathbf{b}_{\mathbf{m}}$ such that

$$\mathbf{s}_k|0\rangle = (P_k + x)|0\rangle.$$

Hence, from (4.4)ii) and (4.3) we have

$$\mathbf{s}_k \in P_k + x + \bigoplus_{\mathbf{m} \notin \{\mathbf{m}(\lambda), \lambda \in \Pi\}} \mathbb{C}_v \mathbf{b}_{\mathbf{m}} \subset P_k + \sum_i E_i \bullet \mathbf{H}_p.$$

Using the induction hypothesis, we obtain

$$\mathbf{s}_k \in (\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \otimes \mathbb{C}_v[\mathbf{l}_1, \dots, \mathbf{l}_r])[k\delta].$$

But $\mathcal{Z} = \mathbb{C}_v[\mathbf{s}_1, \mathbf{s}_2, \dots]$ and hence by Theorem 4.1 the multiplication map

$$(\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_p) \otimes \mathbb{C}_v[\mathbf{l}_1, \dots, \mathbf{l}_r])[k\delta] \rightarrow \mathbf{H}_p[k\delta]$$

is surjective. By graded dimension argument it is also injective. This proves the first assertion of the Lemma. The second is proved in a similar way. $\diamond$

The following result makes the link between $\mathbf{U}_v^+(\widehat{\mathfrak{gl}}_p)$ and $\mathbf{H}_p$ explicit at the classical level.

Lemma 4.2. *There exists a unique isomorphism $\varphi : \mathbf{U}^+(\widehat{\mathfrak{gl}}_p) \xrightarrow{\sim} (\mathbf{H}_p)_{|v=1}$ extending the canonical embedding $\mathbf{U}^-(\widehat{\mathfrak{sl}}_n) \hookrightarrow (\mathbf{H}_p)_{|v=1}$ such that $\varphi(ht^r) = \mathbf{h}_r$ for all $r \geq 1$.*

Proof. It follows from [Ri2] that under the canonical embedding $\mathbf{U}^-(\widehat{\mathfrak{sl}}_n) \hookrightarrow (\mathbf{H}_p)_{|v=1}$ we have, for $i = 1, \dots, n-1$ and $s \geq 0, r \geq 1$

$$e_i t^s = \pm[i; 1 + ps), \quad f_i t^r = \pm[i + 1; pr - 1), \quad h_i t^r = \pm([i; lr) - [i - 1; lr)).$$

A direct computation then shows that the assignement $ht^r \mapsto \mathbf{h}_r$ extends to a well-defined algebra homomorphism. The result now follows from Lemma 4.1. $\diamond$

4.3. Let $\mathbf{U}_v(\widehat{\mathfrak{sl}}_p)$ be the quantized enveloping algebra of the affine Kac-Moody algebra $\widehat{\mathfrak{sl}}_p$ and let $\mathbf{U}_v^{(D)}(\widehat{\mathfrak{sl}}_p)$ be Drinfeld's new realization (see [Be]). In [Be], Beck explicitly constructed an isomorphism of $\mathbb{C}(v)$ -algebras $i : \mathbf{U}_v(\widehat{\mathfrak{sl}}_p) \xrightarrow{\sim} \mathbf{U}_v^{(D)}(\widehat{\mathfrak{sl}}_p)$. Let $\pi : \mathbf{U}_v^{(D)}(\widehat{\mathfrak{sl}}_p) \rightarrow \mathbf{U}_v(L\mathfrak{sl}_p)$ be the canonical projection (obtained by setting the central charge $c = 0$). Set $\epsilon_0 = \pi \circ i(E_0)$ and let $\mathbf{U}_0^+ \subset \mathbf{U}_v(L\mathfrak{sl}_p)$ be the subalgebra generated by ϵ_0 and $x_{i,0}^+$ for $i = 1, \dots, p-1$. Let $\mathbf{U}^+ \subset \mathbf{U}_v(L\mathfrak{gl}_p)$ be the $\mathbb{C}_v$ -subalgebra generated by $\mathbf{U}_0^+$ and $h_{0,l}$ for $l > 0$. The quantum analogue of Lemma 4.2 is the following conjecture :

Conjecture. *The assignment $\epsilon_0 \mapsto E_0, x_{i,0}^+ \mapsto E_i, h_{0,l} \mapsto \mathbf{h}_l$ extends to an isomorphism of $\mathbb{C}(v)$ -algebras $\mathbf{U}^+ \otimes \mathbb{C}(v) \xrightarrow{\sim} \mathbf{H}_p \otimes \mathbb{C}(v)$.*

4.4. Let Γ be a star Dynkin diagram as in Section 1.4. We will now define a quantum analog of the subalgebra $\mathbf{U}(\widehat{\mathfrak{n}}) \subset \mathbf{U}(\widehat{\mathfrak{g}})$. To each type A subdiagram J_s , $s = 1, \dots, n$ we associate the algebra $\mathbf{H}_{p_s}$ where $\phi_s : A_{p_s} \xrightarrow{\sim} J_s$ and we denote by $E_0^{(s)}, E_{\phi_s(1)}, \dots, E_{\phi_s(p_s-1)}$ and $\mathbf{h}_r^{(s)} \in \mathbf{H}_{p_s}$ the corresponding elements, as defined in Section 4.2.

Definition. Let $\mathbf{U}_v(\widehat{\mathfrak{n}})$ be the $\mathbb{C}_v$ -algebra generated by $\mathbf{H}_{p_s}$ for $s = 1, \dots, n$, $H_{\star,r}$, for $r \in \mathbb{N}^*$ and $E_{\star,k}$ for $k \in \mathbb{Z}$, subject to the following set of relations :

- i) For all s , $H_{\star,r} = \mathbf{h}_r^{(s)}$, and $[\mathbf{H}_{p_s}, \mathbf{H}_{p_{s'}}] = 0$ for $s \neq s'$.
- ii) We have

$$(4.7) \quad [H_{\star,r}, E_{\star,t}] = \frac{[2r]}{r} E_{\star,r+t},$$

$$(4.8) \quad E_{\star,t_1+1} E_{\star,t_2} - v^2 E_{\star,t_2} E_{\star,t_1+1} = v^2 E_{\star,t_1} E_{\star,t_2+1} - E_{\star,t_2+1} E_{\star,t_1},$$

iii)

$$(4.9) \quad a_{\star,j} = 0 \Rightarrow [E_{\star,t}, E_j] = 0, \quad v E_{\star,t} E_0^{(s)} - E_0^{(s)} E_{\star,t} = 0,$$

$$(4.10) \quad p_s > 2 \Rightarrow [E_0^{(s)}, v E_{\phi_s(1)} E_{\star,t} - E_{\star,t} E_{\phi_s(1)}] = 0,$$

- iv) For $s = 1, \dots, n$ and $r \geq 1$ set $E_{\phi_s(1)}^{(r)} = \frac{r}{[r]} [E_{\phi_s(1)}, H_{\star,r}]$. Then for any $r, r_1, r_2 \in \mathbb{N}$ and $t, t_1, t_2 \in \mathbb{Z}$,

$$(4.11) \quad \text{Sym}_{r_1, r_2} \{ E_{\phi_s(1)}^{(r_1)} E_{\phi_s(1)}^{(r_2)} E_{\star,t} - [2] E_{\phi_s(1)}^{(r_1)} E_{\star,t} E_{\phi_s(1)}^{(r_2)} + E_{\star,t} E_{\phi_s(1)}^{(r_1)} E_{\phi_s(1)}^{(r_2)} \} = 0,$$

$$(4.12) \quad \text{Sym}_{t_1, t_2} \{ E_{\star,t_1} E_{\star,t_2} E_{\phi_s(1)}^{(r)} - [2] E_{\star,t_1} E_{\phi_s(1)}^{(r)} E_{\star,t_2} + E_{\phi_s(1)}^{(r)} E_{\star,t_1} E_{\star,t_2} \} = 0,$$

- v) For any $s = 1, \dots, n$ and $r, r_1, r_2 \in \mathbb{N}$, $t \in \mathbb{Z}$ we have

$$(4.13) \quad E_{\star,t+1} E_{\phi_s(1)}^{(r)} - v^{-1} E_{\phi_s(1)}^{(r)} E_{\star,t+1} = v^{-1} E_{\star,t} E_{\phi_s(1)}^{(r+1)} - E_{\phi_s(1)}^{(r+1)} E_{\star,t},$$

$$(4.14) \quad E_{\phi_s(1)}^{(r_1+1)} E_{\phi_s(1)}^{(r_2)} - v^2 E_{\phi_s(1)}^{(r_2)} E_{\phi_s(1)}^{(r_1+1)} = v^2 E_{\phi_s(1)}^{(r_2+1)} E_{\phi_s(1)}^{(r_1)} - E_{\phi_s(1)}^{(r_1)} E_{\phi_s(1)}^{(r_2+1)}.$$

Note that $E_{\phi_s(1)}^{(r)} \in \mathbf{H}_{p_s}$ by (4.1). Moreover, if Conjecture 4.3 is true then there is a natural homomorphism $\mathbf{U}_v(\widehat{\mathfrak{n}}) \rightarrow \mathbf{U}_v(L\mathfrak{g})$.

5. MAIN THEOREMS

5.1. Fix a finite field k with q elements and let $\mathbf{p} = (p_1, \dots, p_n) \in \mathbb{N}^n$, $\underline{\lambda} = (\lambda_1, \dots, \lambda_n) \in (\mathbb{P}^1(k))^n$ be as in Section 2.1. Let Γ be the corresponding star Dynkin diagram, and let $\mathfrak{g}, \widehat{\mathfrak{g}}, \widehat{\mathfrak{n}}$, etc... be associated to Γ as in Section 1. Observe that, from Proposition 2.4 there is a natural identification of $\mathbb{Z}$ -modules $K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})) \xrightarrow{\sim} \widehat{Q}$ given by the assignement

$$\begin{aligned} [S_i^{(s)}] &\mapsto \alpha_{\phi_s^{-1}(i)} \text{ for } i = 1, \dots, p_s - 1, \\ [S_0^{(s)}] &\mapsto \delta - \theta^{(s)}, \\ [\mathcal{O}_{\mathbb{X}}(k\vec{c})] &\mapsto \alpha_{\star} + k\delta. \end{aligned}$$

Define the symmetric bilinear form $(,) : K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})) \otimes K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})) \rightarrow \mathbb{Z}$ by $(M, N) = \langle M, N \rangle + \langle N, M \rangle$.

Proposition 5.1. *The bilinear form $(,)$ on $K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}}))$ coincides with the Cartan form on $\widehat{Q}$.*

This is a direct consequence of Lemma 6.1, proved in Section 6.2.

5.2. Let x be a closed point of $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ and let $\mathcal{T}_x$ be the Serre subcategory of $\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$ consisting of (torsion) sheaves supported at x . Note that there is a canonical embedding of Hall algebras

$$\mathbf{H}_{\mathcal{T}_x} \hookrightarrow \mathbf{H}_{\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})}.$$

If x is an ordinary closed point of degree pd then there is a canonical isomorphism

$$\Theta_x : \mathbf{H}_1^{(q^d)} \xrightarrow{\sim} \mathbf{H}_{\mathcal{T}_x}.$$

For every $r \in \mathbb{N}$ we set

$$\mathbf{h}_{r,x} = \begin{cases} 0 & \text{if } r \not\equiv 0 \pmod{d} \\ \Theta_x(\mathbf{h}_{r/d}) & \text{if } r \equiv 0 \pmod{d} \end{cases}$$

where $\mathbf{h}_l \in \mathbf{H}_1^{(q^d)}$ is defined in Section 4.1.

If $x = \sigma_s$ for some $s = 1, \dots, n$ then x is of degree $\frac{p}{p_s}$ and there is a canonical isomorphism

$$\Theta_{\sigma_s} : \mathbf{H}_{p_s}^{(q)} \xrightarrow{\sim} \mathbf{H}_{\mathcal{T}_{\sigma_s}}.$$

For every $r \in \mathbb{N}$ and $s = 1, \dots, n$ we set

$$\mathbf{h}_{r,\sigma_s} = \Theta_{\sigma_s}(\mathbf{h}_r)$$

where $\mathbf{h}_r$ is defined in Section 4.2. Finally we set

$$T_r = \bigoplus_x \mathbf{h}_{r,x} \in \mathbf{H}_{\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})},$$

where the sum ranges over all closed points of $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$. This sum is finite as $\mathbf{h}_{r,x} = 0$ for all but finitely many x .

Recall that $S_i^{(s)}$ for $i = 0, \dots, p_s - 1$ denotes the simple torsion sheaves supported at σ_s (see Section 2.4.).

5.3. Let $\mathbf{U}_{\mathbf{p},\Delta}$ be the subalgebra of $\mathbf{H}_{\text{Coh}(\mathbb{X}_{\mathbf{p},\Delta})}$ generated by $[S_i^{(s)}]$ for $s = 1, \dots, n$ and $i = 0, \dots, p_s - 1$, by $[\mathcal{O}_{\mathbb{X}}(k\vec{c})]$ for $k \in \mathbb{Z}$ and by T_r for $r \in \mathbb{N}^*$. The following is our first main result and will be proved in Section 6.

Theorem 5.1. *The assignment $E_{\phi_s(i)} \mapsto [S_i^{(s)}]$, $E_0^{(s)} \mapsto [S_0^{(s)}]$, $H_{\star,r} \mapsto T_r$, $E_{\star,k} \mapsto [\mathcal{O}_{\mathbb{X}}(k\vec{c})]$ extends to a homomorphism of algebras $\Phi : \mathbf{U}_v(\hat{\mathfrak{n}}) \otimes_{\mathbb{C}_v} \mathbb{C}_q \rightarrow \mathbf{U}_{\mathbf{p},\Delta}$.*

Note that $\mathbf{U}_v(\hat{\mathfrak{n}})$ and $\mathbf{U}_{\mathbf{p},\Delta}$ are both $\hat{Q}$ -graded and that Φ is compatible with this grading.

5.4. Following Ringel, we introduce a “generic” analogue of the Hall algebra $\mathbf{U}_{\mathbf{p},\Delta}$. For each finite field k we denote by $\mathbf{H}^{(k)}$ the Hall algebra of the category of coherent sheaves on the weighted projective line $\mathbb{X}_{\mathbf{p},\Delta}$ defined over k . We will use the notations $[S_i^{(s)}]^{(k)}$, $T_r^{(k)}$, etc... for the corresponding elements. Consider the direct product of rings $\mathbf{H} = \prod_k \mathbf{H}^{(k)}$. Let $\check{\mathbf{U}}_{\mathbf{p},\Delta}$ be the subalgebra of $\mathbf{H}$ generated by the collection of elements

$$\begin{aligned} [S_i^{(s)}] &= \prod_k [S_i^{(s)}]^{(k)}, \\ T_r &= \prod_k T_r^{(k)}, \\ [\mathcal{O}_{\mathbb{X}}(l\vec{c})] &= \prod_k [\mathcal{O}_{\mathbb{X}}(l\vec{c})]^{(k)}. \end{aligned}$$

Both $\mathbf{H}$ and $\check{\mathbf{U}}_{\mathbf{p},\Delta}$ are $\mathbb{C}_v$ -modules since each $\mathbf{H}^{(k)}$ is a $\mathbb{C}_v$ -module. Moreover, the element $v \in \mathbf{H}$ does not satisfy any polynomial equations $P(v) = 0$ and we can consider it as a formal parameter. In particular, the algebra $\check{\mathbf{U}}_{\mathbf{p},\Delta} \otimes_{\mathbb{C}_v} \mathbb{C}(v)$ is well-defined.

Our second main result is the following.

Theorem 5.2. *Assume that Γ is finite or affine. The assignment $E_{\phi_s(i)} \mapsto [S_i^{(s)}]$, $E_0^{(s)} \mapsto [S_0^{(s)}]$, $H_{\star,r} \mapsto T_r$, $E_{\star,k} \mapsto [\mathcal{O}_{\mathbb{X}}(k\vec{c})]$ extends to an isomorphism of algebras*

$$\check{\Phi} : \mathbf{U}_v(\hat{\mathfrak{n}}) \otimes_{\mathbb{C}_v} \mathbb{C}(v) \xrightarrow{\sim} \check{\mathbf{U}}_{\mathbf{p},\Delta} \otimes_{\mathbb{C}_v} \mathbb{C}(v).$$

This theorem is proved in Section 7 (for the finite type case) and in Section 8 (for the affine case).

Remark. We believe that the map Φ in Theorem 5.1 is in fact an isomorphism (for all Γ). When Γ is finite or affine, this would follow from the results of Section 7 and 8 together with the flatness of the deformation $\mathbf{U}_v(\hat{\mathfrak{n}})$ of $\mathbf{U}(\hat{\mathfrak{n}})$.

6. COMPUTATIONS OF SOME HALL NUMBERS

In this section we check that the map Φ is well-defined, i.e that relations (4.7)-(4.14) are satisfied in the Hall algebra. This proves Theorem 5.1. For simplicity we write $\mathbb{X}$ for $\mathbb{X}_{\mathbf{p},\Delta}$ and $\mathcal{O}$ for the structure sheaf $\mathcal{O}_{\mathbb{X}}$.

6.1. It is clear that if S and S' are torsion sheaves with disjoint support then $[S][S'] = [S \oplus S'] = [S'][S]$ in $\mathbf{H}_{\text{Coh}(\mathbb{X})}$. Moreover it follows from Sections 4.1 and 4.2 that the subalgebra generated by T_r for $r \geq 1$ is commutative. Hence relation i) in Section 4.4 is satisfied in $\mathbf{U}_{\mathbf{p},\Delta}$.

6.2. Recall that $\langle , \rangle$ denotes the Euler form on $K_0(\text{Coh}(\mathbb{X}))$. From (2.1) and Theorem 2.5 one easily deduces the following lemma. Set

$$\delta = [\mathcal{O}(\vec{c})] - [\mathcal{O}] = \sum_{i=0}^{p_1-1} [S_i^{(1)}] = \cdots = \sum_{i=0}^{p_s-1} [S_i^{(s)}] \in K_0(\text{Coh}(\mathbb{X})).$$

This corresponds to the imaginary root $\delta \in \widehat{Q}$ under the identification

$$K_0(\text{Coh}(\mathbb{X})) \simeq \widehat{Q}.$$

Lemma 6.1. *The Euler form $\langle , \rangle$ is given by :*

$$\begin{aligned} \langle [\mathcal{O}], [\mathcal{O}] \rangle &= 1, & \langle [\mathcal{O}], \delta \rangle &= 1, & \langle \delta, [\mathcal{O}] \rangle &= -1, \\ \langle \delta, \delta \rangle &= 0, & \langle \delta, [S_i^{(s)}] \rangle &= 0, & \langle [S_i^{(s)}], \delta \rangle &= 0, \end{aligned}$$

and

$$\begin{aligned} \langle \mathcal{O}, [S_i^{(s)}] \rangle &= \begin{cases} 1 & \text{if } i = p_s \\ 0 & \text{if } i \neq p_s \end{cases} \\ \langle [S_i^{(s)}], \mathcal{O} \rangle &= \begin{cases} -1 & \text{if } i = 1 \\ 0 & \text{if } i \neq 1 \end{cases} \\ \langle [S_i^{(s)}], [S_{i'}^{(s')}] \rangle &= \begin{cases} 1 & \text{if } s = s', i = i' \\ -1 & \text{if } s = s', i \equiv i' + 1 \pmod{p_s} \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

6.3. Let $\mathcal{C}_0$ be the Serre subcategory of $\text{Coh}(\mathbb{X})$ generated by the locally free sheaves $\mathcal{O}(k\vec{c})$ for $k \in \mathbb{N}$, by the torsion sheaves supported at ordinary closed points, and by the sheaves $S_{r\delta}^{(s)}$ for $s = 1, \dots, n$ and $r \geq 1$ defined by the exact sequences

$$0 \rightarrow \mathcal{O} \rightarrow \mathcal{O}(r\vec{c}) \rightarrow S_{r\delta}^{(s)} \rightarrow 0$$

induced by multiplication by $x_s^{rp_s}$. Let $\text{Coh}(\mathbb{P}^1(k))$ be the category of coherent sheaves on $\mathbb{P}^1(k)$.

Lemma 6.2. *The categories $\mathcal{C}_0$ and $\text{Coh}(\mathbb{P}^1(k))$ are equivalent.*

Proof. By Serre's theorem and by Proposition 2.3, there is a chain of equivalences

$$\begin{aligned} \text{Coh}(\mathbb{P}^1(k)) &\simeq k[T, U] - \text{modgr} / k[T, U] - \text{modgr}_0 \\ &\simeq S(\mathbf{p}, \underline{\lambda})^0 - \text{modgr} / S(\mathbf{p}, \underline{\lambda})^0 - \text{modgr}_0 \\ &\simeq \mathcal{C}_0 \end{aligned}$$

where $S(\mathbf{p}, \underline{\lambda})^0$ denotes the $(\mathbb{Z}$ -graded) subring of $S(\mathbf{p}, \underline{\lambda})$ consisting of elements of degree $k\vec{c}$ for some $k \geq 0$. $\diamond$

The structure of the Hall algebra of $\text{Coh}(\mathbb{P}^1(k))$ has been unraveled by Kapranov [Kap] and recovered by Baumann and Kassel in [BK]. In particular, relation (4.8) now follows from Lemma 6.2 and [BK], Lemma 16 i).

For every ordinary closed point x of degree pd we set

$$\Xi_x(s) = 1 + \sum_{\beta \in \text{Iso}T_x} [\beta] s^{\mathbf{d}(\beta)/p} = 1 + \sum_{r \geq 1} \Theta_x \circ \Upsilon(\xi_r) s^{rd},$$

where $\xi_r \in \Gamma$ denotes the complete symmetric function (we avoid the usual notation h_r by fear of confusion with $\mathbf{h}_r$). For every exceptional point σ_i we set

$$\Xi_{\sigma_i}(s) = 1 + \sum_{r \geq 1} \Theta_{\sigma_i} \circ \Psi \circ \Upsilon(\xi_r) s^r.$$

Now define $\widehat{\xi}_r \in \mathbf{H}_{Coh(\mathbb{X})}$ by the relation

$$1 + \sum_{r \geq 1} \widehat{\xi}_r s^r = \prod_{x \in \mathbb{X}_{\mathbf{p}, \Delta}} \Xi_x(s).$$

Lemma 6.3. *We have*

$$1 + \sum_{r \geq 1} \widehat{\xi}_r s^r = \exp\left(\sum_{r \geq 1} \frac{T_r}{[r]} s^r\right).$$

Proof. Classes of torsion sheaves supported at distinct points of $\mathbb{X}$ commute in the Hall algebra $\mathbf{H}_{Coh(\mathbb{X})}$. Hence,

$$\exp\left(\sum_{r \geq 1} \frac{T_r}{[r]} s^r\right) = \prod_{x \in \mathbb{X}} \exp\left(\sum_{r \geq 1} \frac{\mathbf{h}_{r,x}}{[r]}\right).$$

Thus it is enough to show that for any closed point x we have

$$\Xi_x(s) = \exp\left(\sum_{r \geq 1} \frac{\mathbf{h}_{r,x}}{[r]}\right).$$

This is a consequence of the definitions of Ξ_r and $\mathbf{h}_{r,x}$ and of the following identity in the ring Γ (see [M], I.2) :

$$\sum_{r \geq 0} \xi_r s^r = \exp\left(\sum_{r \geq 1} \frac{p_r}{r}\right).$$

◇

From Lemma 6.2 and [BK], Lemma 19 we have, for any $r \geq 1$ and any $n \in \mathbb{Z}$,

$$(6.1) \quad \widehat{\xi}_r[\mathcal{O}(n\vec{c})] = [r+1][\mathcal{O}((n+r)\vec{c})] + \sum_{s=0}^{r-1} [\mathcal{O}((n+s)\vec{c})] \widehat{\xi}_{r-s}.$$

Relation (4.7) is now a consequence of Lemma 6.3 and (6.1) (see [BK], Proposition 25).

6.4. We now check relations (4.9) and (4.10). Note that the group $L(\mathbf{p})$ acts on $\mathbf{H}_{Coh(\mathbb{X})}$. Thus from (2.2) we can assume $t = 0$. If $a_{*,j} = 0$ and $j = \phi_s(k)$ we have, using Serre duality,

$$\mathrm{Ext}^1(S_k^{(s)}, \mathcal{O}) = \mathrm{Ext}^1(\mathcal{O}, S_k^{(s)}) = 0$$

and

$$\mathrm{Hom}(\mathcal{O}, S_k^{(s)}) = \mathrm{Ext}^1(S_k^{(s)}, \mathcal{O}(\vec{\omega})) = \mathrm{Ext}^1(S_{k+1}^{(s)}, \mathcal{O}) = 0.$$

Moreover, $\langle [\mathcal{O}], [S_k^{(s)}] \rangle = \langle [S_k^{(s)}], [\mathcal{O}] \rangle = 0$. The first equality in (4.9) follows. Similarly, we have

$$\mathrm{Ext}^1(S_{p_s}^{(s)}, \mathcal{O}) = \mathrm{Ext}^1(\mathcal{O}, S_{p_s}^{(s)}) = 0, \quad \mathrm{Hom}(\mathcal{O}, S_{p_s}^{(s)}) = k.$$

Thus,

$$\begin{aligned} [S_{p_s}^{(s)}] \cdot [\mathcal{O}] &= v^2[\mathcal{O} \oplus S_{p_s}^{(s)}] \\ [\mathcal{O}] \cdot [S_{p_s}^{(s)}] &= v[\mathcal{O} \oplus S_{p_s}^{(s)}] \end{aligned}$$

and the second equality of (4.9) follows. To prove (4.10) we compute in an analogous fashion

$$\begin{aligned} [S_1^{(s)}] \cdot [\mathcal{O}] &= v^{-1}([\mathcal{O} \oplus S_1^{(s)}] + [\mathcal{O}(\vec{x}_i)]) \\ [\mathcal{O}] \cdot [S_1^{(s)}] &= [\mathcal{O} \oplus S_1^{(s)}]. \end{aligned}$$

Hence,

$$(6.2) \quad v[S_1^{(s)}] \cdot [\mathcal{O}] - [\mathcal{O}] \cdot [S_1^{(s)}] = [\mathcal{O}(\vec{x}_i)].$$

If $p_s > 2$ then from (4.9) we deduce

$$[[S_{p_s}^{(s)}], [\mathcal{O}(\vec{x}_i)]] = [[S_{p_s-1}^{(s)}], [\mathcal{O}]](\vec{x}_i) = 0$$

which proves (4.10).

6.5. Let us set $[S_{1,r}^{(s)}] = \frac{r}{[r]} [[S_1^{(s)}], T_r]$. An explicit computation in $\mathbf{H}_{p_s}$ shows that

$$[S_{1,r}^{(s)}] = \sum_{\lambda, |\lambda|=r} n(l(\lambda)) [|\lambda\rangle^{(s)} \oplus S_1^{(s)}] + \sum_{\lambda, |\lambda|=r} n(l(\lambda) - 1) \sum_{\nu \triangleleft \lambda} [[1; p_s \nu + 1] \oplus |\lambda \setminus \nu\rangle^{(s)}],$$

where for simplicity we set $|\lambda\rangle^{(s)} = \Theta_{\sigma_s}(f_\lambda)$ and where we write $\nu \triangleleft \lambda$ if $\nu \in \mathbb{N}$ is a part of λ . The following lemma will be important for us.

Lemma 6.4. *For any $r_1, r_2 \in \mathbb{N}$ we have*

$$(6.3) \quad [[S_{1,r_2}^{(s)}], T_{r_1}] = \frac{[r_1]}{r_1} [S_{1,r_1+r_2}^{(s)}].$$

Proof. This is an explicit computation in $\mathbf{H}_p$, and is proved in Appendix 1.

6.6. In this section we check relation (4.11). From (6.2) and the relation $v[\mathcal{O}(\vec{x}_i)] \cdot [S_1^{(s)}] - [S_1^{(s)}] \cdot [\mathcal{O}(\vec{x}_i)] = 0$ we deduce

$$\begin{aligned} 0 &= v([S_1^{(s)}] \cdot [\mathcal{O}] - [\mathcal{O}] \cdot [S_1^{(s)}]) \cdot [S_1^{(s)}] - [S_1^{(s)}] \cdot (v[S_1^{(s)}] \cdot [\mathcal{O}] - [\mathcal{O}] \cdot [S_1^{(s)}]) \\ &= [S_1^{(s)}]^2 \cdot [\mathcal{O}] - [2][S_1^{(s)}] \cdot [\mathcal{O}] \cdot [S_1^{(s)}] + [\mathcal{O}] \cdot [S_1^{(s)}]^2 \end{aligned}$$

proving (4.11) when $r_1 = r_2 = t = 0$. By twisting by $t\vec{c}$ also obtain the case $r_1 = r_2 = 0$ and $t \in \mathbb{Z}$. Now let us apply *ad* T_{r_1} to (6). We get, using (4.7)

$$\begin{aligned} 0 &= \frac{[r_1]}{r_1} \text{Sym}_{r_1,0} \{ [S_{1,r_1}^{(s)}] \cdot [S_{1,0}^{(s)}] \cdot [\mathcal{O}] - [2][S_{1,r_1}^{(s)}] \cdot [\mathcal{O}] \cdot [S_{1,0}^{(s)}] + [\mathcal{O}] \cdot [S_{1,r_1}^{(s)}] \cdot [S_{1,0}^{(s)}] \} \\ &\quad - \frac{[2r_1]}{r_1} \{ [S_1^{(s)}]^2 \cdot [\mathcal{O}(r_1\vec{c})] - [2][S_1^{(s)}] \cdot [\mathcal{O}(r_1\vec{c})] \cdot [S_1^{(s)}] + [\mathcal{O}(r_1\vec{c})] \cdot [S_1^{(s)}]^2 \}, \end{aligned}$$

Hence,

$$(6.4) \quad \text{Sym}_{r_1,0} \{ [S_{1,r_1}^{(s)}] \cdot [S_{1,0}^{(s)}] \cdot [\mathcal{O}] - [2][S_{1,r_1}^{(s)}] \cdot [\mathcal{O}] \cdot [S_{1,0}^{(s)}] + [\mathcal{O}] \cdot [S_{1,r_1}^{(s)}] \cdot [S_{1,0}^{(s)}] \} = 0,$$

proving (4.11) when $r_2 = 0$. Finally, we apply *ad* T_{r_2} to (6.4). Using Lemma 6.4, we get

$$\begin{aligned} 0 &= -\frac{[r_2]}{r_2} \text{Sym}_{r_1+r_2,0} \{ [S_{1,r_1+r_2}^{(s)}] \cdot [S_{1,0}^{(s)}] \cdot [\mathcal{O}] - [2][S_{1,r_1+r_2}^{(s)}] \cdot [\mathcal{O}] \cdot [S_{1,0}^{(s)}] \\ &\quad + [\mathcal{O}] \cdot [S_{1,r_1+r_2}^{(s)}] \cdot [S_{1,0}^{(s)}] \} \\ &\quad - \frac{[r_2]}{r_2} \text{Sym}_{r_1,r_2} \{ [S_{1,r_1}^{(s)}] \cdot [S_{1,r_2}^{(s)}] \cdot [\mathcal{O}] - [2][S_{1,r_1}^{(s)}] \cdot [\mathcal{O}] \cdot [S_{1,r_2}^{(s)}] \\ &\quad + [\mathcal{O}] \cdot [S_{1,r_1}^{(s)}] \cdot [S_{1,r_2}^{(s)}] \} \\ &\quad + \frac{[2r_2]}{r_2} \text{Sym}_{r_1,0} \{ [S_{1,r_1}^{(s)}] \cdot [S_{1,0}^{(s)}] \cdot [\mathcal{O}(r_2\vec{c})] - [2][S_{1,r_1}^{(s)}] \cdot [\mathcal{O}(r_2\vec{c})] \cdot [S_{1,0}^{(s)}] \\ &\quad + [\mathcal{O}(r_2\vec{c})] \cdot [S_{1,r_1}^{(s)}] \cdot [S_{1,0}^{(s)}] \}. \end{aligned}$$

Using (6.4) (with $r_1 + r_2$) we deduce that (4.11) holds for all r_1, r_2 and t .

6.7. We now prove relation (4.12). Reasoning as in Section 6.6 and using Lemma 6.5 we see that it is enough to deal with the case $t_2 = r = 0$. Twisting (6.2) by $n\vec{c}$ we obtain

$$(6.5) \quad v[S_1^{(s)}] \cdot [\mathcal{O}(t\vec{c})] - [\mathcal{O}(t\vec{c})] \cdot [S_1^{(s)}] = [\mathcal{O}(t\vec{c} + \vec{x}_s)].$$

Hence relation (4.12) is equivalent to the relation

$$(6.6) \quad \begin{aligned} 0 &= [\mathcal{O}(t\vec{c})] \cdot [\mathcal{O}(\vec{x}_s)] - v^{-1}[\mathcal{O}(\vec{x}_s)] \cdot [\mathcal{O}(t\vec{c})] \\ &\quad + [\mathcal{O}] \cdot [\mathcal{O}(t\vec{c} + \vec{x}_s)] - v^{-1}[\mathcal{O}(t\vec{c} + \vec{x}_s)] \cdot [\mathcal{O}]. \end{aligned}$$

An explicit computation using Lemma 6.1 gives

$$(6.7) \quad [\mathcal{O}(\vec{x}_s)] \cdot [\mathcal{O}(t\vec{c})] = v^t[\mathcal{O}(t\vec{c}) \oplus \mathcal{O}(\vec{x}_s)],$$

$$(6.8) \quad [\mathcal{O}] \cdot [\mathcal{O}(t\vec{c} + \vec{x}_s)] = v^{1+t}[\mathcal{O} \oplus \mathcal{O}(t\vec{c} + \vec{x}_s)].$$

The computation of the other two products is given by the following lemma. If $i, j \geq 1$ let us denote by $P_{i,j}$ the set of pairs of homogeneous polynomials $(P, Q) \in k[T, U]$ of degrees i and j respectively such that P and Q are relatively prime and such that T does not divide P .

Lemma 6.5. *We have*

$$\begin{aligned} [\mathcal{O}(t\vec{c})] \cdot [\mathcal{O}(\vec{x}_s)] &= v^{1-t} \{ v^{2t}[\mathcal{O}(\vec{x}_s) \oplus \mathcal{O}(t\vec{c})] \\ &\quad + \sum_{k=1}^{t-1} \frac{1}{q-1} |P_{k,t-k-1}| [\mathcal{O}(k\vec{c} + \vec{x}_s) \oplus \mathcal{O}((t-k)\vec{c})] \}, \\ [\mathcal{O}(t\vec{c} + \vec{x}_s)] \cdot [\mathcal{O}] &= v^{-t} \{ v^{2t+2}[\mathcal{O} \oplus \mathcal{O}(t\vec{c} + \vec{x}_s)] + (v^2 - 1)v^{2t}[\mathcal{O}(\vec{x}_s) \oplus \mathcal{O}(t\vec{c})] \\ &\quad + \sum_{k=1}^{t-1} \frac{1}{q-1} |P_{t-k,k}| [\mathcal{O}(k\vec{c} + \vec{x}_s) \oplus \mathcal{O}((t-k)\vec{c})] \}. \end{aligned}$$

Proof. A reasoning similar to that of [BK], Proposition 5, shows that the coefficient $c_{\vec{l}_1, \vec{l}_2}$ of $[\mathcal{O}(\vec{l}_1) \oplus \mathcal{O}(\vec{l}_2)]$ in the product $[\mathcal{O}(t\vec{c})] \cdot [\mathcal{O}(\vec{x}_s)]$ is equal to $\frac{1}{q-1} |Q_{\vec{l}_1, \vec{l}_2}|$ where $Q_{\vec{l}_1, \vec{l}_2}$ is the set of pairs of relatively prime elements $P, Q \in S(\mathbf{p}, \underline{\lambda})$ of respective degrees $\vec{l}_1 - \vec{x}_s$ and $\vec{l}_2 - \vec{x}_s$. Comparing images in the Grothendieck group, it is clear that $c_{\vec{l}_1, \vec{l}_2} = 0$ if $\vec{l}_1 + \vec{l}_2 \neq t\vec{c} + \vec{x}_s$. Moreover, observe that if $\vec{l} = k\vec{c} + t_1\vec{x}_1 + \dots + t_n\vec{x}_n$ with $0 \leq t_i < p_i$ then

$$S(\mathbf{p}, \underline{\lambda})[\vec{l}] = x_1^{t_1} \dots x_n^{t_n} S(\mathbf{p}, \underline{\lambda})[k\vec{c}].$$

From there one easily deduces that $c_{\vec{l}_1, \vec{l}_2} = 0$ unless $\{\vec{l}_1, \vec{l}_2\} = \{k\vec{c} + \vec{x}_s, (n-k)\vec{c}\}$ for some $0 \leq k \leq t-1$. Finally, $S(\mathbf{p}, \underline{\lambda})[k\vec{c}] = k[x_s^{p_s}, x_r^{p_r}]$ for any $r = 1, \dots, n$, $r \neq s$. This provides a bijection between $P_{k,t-k-1}$ and $Q_{k\vec{c}, (t-k)\vec{c} - \vec{x}_s}$ and proves the first relation in the Lemma. The second relation is proved in a similar manner. $\diamond$

From (6.7) and Lemma 6.5 we see that (4.12) is equivalent to the relation

$$\sum_{k=1}^{t-1} (v|P_{k,t-k-1}| - v^{-1}|P_{t-k,k}|) [\mathcal{O}(k\vec{c} + \vec{x}_s) \oplus \mathcal{O}((t-k)\vec{c})] = 0,$$

which in turn is a consequence of the following lemma :

Lemma 6.6. *If $1 \leq k \leq t-1$ then $P_{t-k,k} = v^2 P_{k,n-k-1}$.*

Proof. Let us denote for simplicity by $S_d = (k[T, U])[t]$ the space of homogeneous polynomials of degree d , and let us write $(P, Q) = 1$ if P and Q are relatively prime.

Then

$$\begin{aligned} |P_{d_1, d_2}| &= |\{(P, Q) \in S_{d_1} \times S_{d_2} \mid (P, Q) = 1, (P, T) = 1\}| \\ &= |\{(P, Q) \in S_{d_1} \times S_{d_2} \mid (P, Q) = 1\}| \\ &\quad - |\{(P, Q) \in S_{d_1} \times S_{d_2} \mid (P, Q) = 1, P = TP'\}|. \end{aligned}$$

In a similar way,

$$\begin{aligned} &|\{(P, Q) \in S_{d_1} \times S_{d_2} \mid (P, Q) = 1, P = TP'\}| \\ &= |\{(P', Q) \in S_{d_1-1} \times S_{d_2} \mid (P', Q) = 1, (T, Q) = 1\}| \\ &= P_{d_2, d_1-1}. \end{aligned}$$

From [BK], Lemma 9, we have

$$|\{(P, Q) \in S_{d_1} \times S_{d_2} \mid (P, Q) = 1\}| = (q^2 - 1)(q - 1)q^{d_1+d_2-1}$$

if $d_1, d_2 \geq 1$. Hence if $d_1, d_2 \geq 1$ then

$$(6.9) \quad |P_{d_1, d_2}| = (q^2 - 1)(q - 1)q^{d_1+d_2-1} - |P_{d_2, d_1-1}|.$$

The Lemma now easily follows from (6.9) by induction on $\min(d_1, d_2)$. $\diamond$

6.8. We here prove relation (4.13). Using the action of $L(\mathbf{p})$ and Lemma 6.4 again we see that it is enough to consider the case $r = t = 0$. Using the notations of Lemma 6.4 we have

$$[S_{1,1}^{(s)}] = [[1; p_s + 1]] + (1 - v^2)[|1\rangle^{(s)} \oplus S_1^{(s)}].$$

Relation (4.13) can now be checked through the following explicit computations :

$$[\mathcal{O}] \cdot [[1; p_s + 1]] = v[\mathcal{O} \oplus [1; p_s + 1]]$$

$$[\mathcal{O}] \cdot [|1\rangle^{(s)} \oplus S_1^{(s)}] = v[\mathcal{O} \oplus |1\rangle^{(s)} \oplus S_1^{(s)}]$$

$$[\mathcal{O}(\vec{c})] \cdot [S_1^{(s)}] = [\mathcal{O}(\vec{c}) \oplus S_1^{(s)}]$$

$$[S_1^{(s)}] \cdot [\mathcal{O}(\vec{c})] = v^{-1}[\mathcal{O}(\vec{c}) \oplus S_1^{(s)}] + v^{-1}[\mathcal{O}(\vec{c} + \vec{x}_s)]$$

$$[[1; p_s + 1]] \cdot [\mathcal{O}] = [\mathcal{O} \oplus [1; p_s + 1]] + (1 - v^{-2})[\mathcal{O}(\vec{x}_s) + |1\rangle^{(s)}] + v^{-2}[\mathcal{O}(\vec{c} + \vec{x}_s)]$$

$$[|1\rangle^{(s)} \oplus S_1^{(s)}] \cdot [\mathcal{O}] = [\mathcal{O} \oplus |1\rangle^{(s)} \oplus S_1^{(s)}] + v^{-2}[\mathcal{O}(\vec{x}_s) \oplus |1\rangle^{(s)}] + v^{-2}[\mathcal{O}(\vec{c}) \oplus S_1^{(s)}].$$

$\diamond$

6.9. The last relation (4.14) is also proved via a direct computation in $\mathbf{H}_{p_s}$ similar to Lemma 6.4 (again, it is enough to assume $r_2 = 0$), (see Appendix 1).

7. THE FINITE TYPE CASE

In this section we assume that Γ is finite and prove Theorem 5.2, i.e we prove that the map $\check{\Phi}$ in Theorem 5.2 is injective.

7.1. Observe that the algebra $\mathbf{U}(\widehat{\mathfrak{n}})$ does not have finite-dimensional weight spaces. We introduce an exhaustive filtration as follows. For each $m \in \mathbb{Z}$, let $\widehat{\mathfrak{n}}_{\geq m}$ be the Lie subalgebra of $\widehat{\mathfrak{n}}$ generated by $\widehat{\mathfrak{n}}_s$ for $s = 1, \dots, n$, $h_{\star, r}$ for $r \geq 1$ and $e_{\star, k}$ for $k \geq m$. We put $\mathbf{U}^{\geq m}(\widehat{\mathfrak{n}}) = \mathbf{U}(\widehat{\mathfrak{n}}_{\geq m})$. It is clear that $\mathbf{U}^{\geq m+1}(\widehat{\mathfrak{n}}) \subset \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})$ and that $\mathbf{U}(\widehat{\mathfrak{n}}) = \bigcup_m \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})$. Note that the assignment $e_{\star, t} \mapsto e_{\star, t+1}$ and the identity on all other generators induces an automorphism σ of $\mathbf{U}(\widehat{\mathfrak{n}})$ which restricts to isomorphisms $\sigma : \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}}) \xrightarrow{\sim} \mathbf{U}^{\geq m+1}(\widehat{\mathfrak{n}})$. In particular, for any $\alpha' \in \mathbb{Z}\delta \oplus \bigoplus_{j \neq \star} \mathbb{Z}\alpha_j$, we have

$$(7.1) \quad \dim \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})[\alpha' + d\alpha_{\star}] = \dim \mathbf{U}^{\geq m+1}(\widehat{\mathfrak{n}})[\alpha' + d\alpha_{\star} + d\delta].$$

Lemma 7.1. *For any $m \in \mathbb{Z}$ the algebra $\mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})$ has finite-dimensional weight spaces.*

Proof. By (7.1) it is enough to assume $m = 0$. Recall the notations of Section 1.4. We have $\widehat{\mathfrak{n}}_{\geq 0} = \widehat{\mathfrak{t}} \oplus \widehat{\mathfrak{f}}_{\geq 0}$ where $\widehat{\mathfrak{f}}_{\geq 0} = \mathfrak{l}[t]$. Hence, by the Poincaré-Birkhoff-Witt theorem, we have

$$\mathbf{U}^{\geq 0}(\widehat{\mathfrak{n}}) = \mathbf{U}(\widehat{\mathfrak{t}}) \otimes \mathbf{U}(\widehat{\mathfrak{f}}_{\geq 0}).$$

Both $\mathbf{U}(\widehat{\mathfrak{t}})$ and $\mathbf{U}(\widehat{\mathfrak{f}}_{\geq 0})$ have finite-dimensional weight spaces. Now, if α is a root of $\widehat{\mathfrak{n}}_{\geq 0}$ then $\alpha = \alpha' + d\delta$ with $\alpha' \in \bigoplus_i \mathbb{Z}\alpha_i$ and $d \geq 0$. Note that there are only finitely many weights β of $\widehat{\mathfrak{t}}$ with $\beta = \beta' + d'\delta$ and $d' \leq d$. Hence,

$$\dim \mathbf{U}^{\geq 0}(\widehat{\mathfrak{n}})[\alpha] = \sum_{\substack{\beta = \beta' + d'\delta \\ d' \leq d}} \dim \mathbf{U}(\widehat{\mathfrak{t}})[\beta] \dim \mathbf{U}(\widehat{\mathfrak{f}}_{\geq 0})[\alpha - \beta] < \infty.$$

◇

In a similar manner, we consider the filtration of $\mathbf{U}_v(\widehat{\mathfrak{n}})$ (resp. of $\mathbf{U}_{\mathbf{p}, \lambda}^{(k)}$) by subalgebras $\mathbf{U}_v^{\geq m}(\widehat{\mathfrak{n}})$ (resp. $\mathbf{U}_{\mathbf{p}, \lambda}^{\geq m, (k)}$) generated by $\mathbf{H}_{p_i}$ for $i = 1, \dots, n$ and by $E_{\star, t}$ for $t \geq m$ (resp. generated by the classes of simple torsion sheaves $[S_i^{(s)}]$, by T_r for $r \geq 1$ and by $[\mathcal{O}(t\vec{c})]$ for $t \geq m$).

Let us view $\mathbb{C}$ as $\mathbb{C}_v$ -module via the evaluation $v \mapsto 1$. It follows from Proposition 1.1, Theorem 4.2 and Lemma 4.2 that

$$\mathbf{U}_v(\widehat{\mathfrak{n}}) \otimes_{\mathbb{C}_v} \mathbb{C} \simeq \mathbf{U}(\widehat{\mathfrak{n}}),$$

and that for any $m \in \mathbb{Z}$,

$$\mathbf{U}_v^{\geq m}(\widehat{\mathfrak{n}}) \otimes_{\mathbb{C}_v} \mathbb{C} \simeq \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}}).$$

Hence, for any $\alpha \in \widehat{Q}$ we have

$$(7.2) \quad \dim_{\mathbb{C}(v)} \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})[\alpha] \otimes_{\mathbb{C}_v} \mathbb{C}(v) \leq \dim_{\mathbb{C}} \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})[\alpha].$$

For any k the map $\Phi^{(k)} : \mathbf{U}_v(\widehat{\mathfrak{n}}) \rightarrow \mathbf{U}_{\mathbf{p}, \lambda}^{(k)}$ restricts to a map $\Phi^{(k)} : \mathbf{U}_v^{\geq m}(\widehat{\mathfrak{n}}) \rightarrow \mathbf{U}_{\mathbf{p}, \lambda}^{\geq m, (k)}$ and, varying k , we thus obtain a morphism

$$(7.3) \quad \check{\Phi} : \mathbf{U}_v^{\geq m}(\widehat{\mathfrak{n}}) \otimes_{\mathbb{C}_v} \mathbb{C}(v) \rightarrow \check{\mathbf{U}}_{\mathbf{p}, \lambda}^{\geq m} \otimes_{\mathbb{C}_v} \mathbb{C}(v).$$

Combining (7.2) with (7.3), we see that the injectivity of $\check{\Phi}$ will follow from the system of inequalities (for all $\alpha \in \widehat{Q}$ and all $m \in \mathbb{Z}$)

$$\dim \check{\mathbf{U}}_{\mathbf{p}, \lambda}^{\geq m}[\alpha] \otimes_{\mathbb{C}_v} \mathbb{C}(v) \geq \dim_{\mathbb{C}} \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})[\alpha],$$

which in turn will follow from the system of inequalities

$$(7.4) \quad \dim_{\mathbb{C}} \mathbf{U}_{\mathbf{p}, \lambda}^{\geq m, (k)}[\alpha] \geq \dim_{\mathbb{C}} \mathbf{U}^{\geq m}(\widehat{\mathfrak{n}})[\alpha]$$

for all finite fields k . Finally, observe that the twist by $\vec{c}$ induces isomorphisms $\mathbf{U}_{\mathbf{p},\underline{\lambda}}^{\geq m,(k)} \xrightarrow{\sim} \mathbf{U}_{\mathbf{p},\underline{\lambda}}^{\geq m+1,(k)}$. In particular, for each $\alpha' \in \mathbb{Z}\delta \oplus \bigoplus_{j \neq \star} \mathbb{Z}\alpha_j$, we have

$$(7.5) \quad \dim \mathbf{U}_{\mathbf{p},\underline{\lambda}}^{\geq m,(k)}[\alpha' + d\alpha_{\star}] = \dim \mathbf{U}_{\mathbf{p},\underline{\lambda}}^{\geq m+1,(k)}[\alpha' + d\alpha_{\star} + d\delta].$$

Hence, in view of (7.1), the system of inequalities (7.4) for all $m \in \mathbb{Z}$ follow from the same system for $m = 0$. This set of inequalities (and therefore Theorem 5.2) are proved in the rest of this section.

7.2. From now on we fix a finite field k and drop the superscript (k) in the notation. Consider the sheaf

$$\mathbf{T} = \bigoplus_{0 \leq \vec{x} \leq \vec{c}} \mathcal{O}(\vec{x}).$$

It is proved in [GL] that $\mathbf{T}$ is a tilting sheaf (in the sense of, e.g. [Hap1]), and that $\text{End}(\mathbf{T})$ is a canonical algebra $\Lambda(\mathbf{p})$ of type $(\mathbf{p})$ (see [GL]), which is itself a tilted algebra of a tame hereditary algebra Σ of (extended) Dynkin type $\hat{\Gamma}$. In particular, there is a chain of equivalences

$$D^b(\text{Coh}(\mathbb{X})) \simeq D^b(\text{mod}(\Lambda^{op})) \simeq D^b(\text{mod}(\Sigma))$$

of (bounded) derived categories. This allows one to translate the well-known classification of indecomposable objects in $D^b(\text{mod}(\Sigma))$ and $\text{mod}(\Sigma)$ into the classification of indecomposable sheaves in $\text{Coh}(\mathbb{X})$, as was done in [Le], §5.8 (see also [LM], 5.4.1.)

Proposition 7.1 ([Le]). *The map $\mathcal{F} \rightarrow [\mathcal{F}] \in K_0(\text{coh}(\mathbb{X})) = \hat{Q}$ induces a bijection between the set of (isomorphism classes) of indecomposable vector bundles and the root system of $\hat{\mathfrak{f}}$.*

We begin with the following lemma.

Lemma 7.2. *The algebra $\mathbf{U}_{\mathbf{p},\underline{\lambda}}$ contains the classes $[\mathcal{O}(\vec{x})]$ of all the line bundles in $\text{Coh}(\mathbb{X}_{\mathbf{p},\underline{\lambda}})$. Moreover, the subalgebra $\mathbf{U}_{\mathbf{p},\underline{\lambda}}^{\geq 0}$ contains the class of all line bundles of the form $\mathcal{O}(\vec{x})$ with $\vec{x} \geq 0$.*

Proof. From (2.2), the set of generators of $\mathbf{U}_{\mathbf{p},\underline{\lambda}}$ is closed under twisting by $k\vec{c} \in L(\mathbf{p})$ for $k \in \mathbb{Z}$. Thus the set of $\vec{x} \in L(\mathbf{p})$ such that $[\mathcal{O}(\vec{x})] \in \mathbf{U}_{\mathbf{p},\underline{\lambda}}$ is closed under twisting by $k\vec{c}$. But from (6.2) we deduce that this set is also invariant under twisting by $\vec{x}_i$ for $i = 1, \dots, n$. Hence, $[\mathcal{O}(\vec{x})] \in \mathbf{U}_{\mathbf{p},\underline{\lambda}}$ for all $\vec{x} \in L(\mathbf{p})$ as desired. $\diamond$

Proposition 7.2. *The algebra $\mathbf{U}_{\mathbf{p},\underline{\lambda}}$ contains the classes of all indecomposable locally free sheaves.*

Proof. By Lemma 7.2, $\mathbf{U}_{\mathbf{p},\underline{\lambda}}$ contains all line bundles. We argue by induction on the rank. Assume that $\mathbf{U}_{\mathbf{p},\underline{\lambda}}$ contains all indecomposable vector bundles of rank r , and let $\mathcal{F}$ be an indecomposable vector bundle of rank $r+1$. By [GL], Proposition 2.6, there exists a line bundle $\mathcal{L}$, a vector bundle $\mathcal{G}$ of rank r and an exact sequence

$$0 \rightarrow \mathcal{L} \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow 0.$$

Let us decompose $\mathcal{G} = \mathcal{G}_1 \oplus \dots \oplus \mathcal{G}_s$ into indecomposable vector bundles, ordered such that $\mu(\mathcal{G}_1) \leq \mu(\mathcal{G}_2) \leq \dots \leq \mu(\mathcal{G}_s)$. By assumption, $[\mathcal{G}_i] \in \mathbf{U}_{\mathbf{p},\underline{\lambda}}$ for each i . Furthermore, from Proposition 2.5 and Corollary 2.1 it follows that $\text{Ext}^1(\mathcal{G}_i, \mathcal{G}_j) = 0$ if $i \leq j$. Thus

$$c_{\mathcal{G}}[\mathcal{G}] = [\mathcal{G}_1] \cdot [\mathcal{G}_2] \cdots [\mathcal{G}_s] \in \mathbf{U}_{\mathbf{p},\underline{\lambda}}$$

for some nonzero $c_{\mathcal{G}} \in \mathbb{C}_v$, and $[\mathcal{G}] \in \mathbf{U}_{\mathbf{p},\underline{\lambda}}$. Now let us write

$$[\mathcal{L}] \cdot [\mathcal{G}] = c_{\mathcal{F}}[\mathcal{F}] + R, \quad \text{for some } 0 \neq c_{\mathcal{F}} \in \mathbb{C}_v.$$

Since there is at most one indecomposable vector bundle with a given class in $K_0(\text{Coh}(\mathbb{X}))$ it follows that R is a sum of terms of the form $c_{\mathcal{H}}[\mathcal{H}]$ where H is a direct sum of indecomposable sheaves of rank at most r . By the same reasoning as for $\mathcal{G}$ we conclude that $[\mathcal{H}] \in \mathbf{U}_{\mathbf{p}, \Delta}$. Thus $[\mathcal{F}] \in \mathbf{U}_{\mathbf{p}, \Delta}$ and the induction step is complete. $\diamond$

7.3. In order to prove (7.4), we need a refinement of Proposition 7.1 and Proposition 7.2. The following Lemma is proved in exactly the same manner as Lemma 7.2.

Lemma 7.3. *The subalgebra $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ contains the classes of all line bundles of the form $\mathcal{O}(\vec{x})$ with $\vec{x} \geq 0$.*

$\diamond$

Now let us consider the full subcategories

$$X_0 = \{\mathcal{F} \in \text{Coh}(\mathbb{X}) \mid \text{Ext}^1(\mathbf{T}, \mathcal{F}) = 0\},$$

$$X_1 = \{\mathcal{F} \in \text{Coh}(\mathbb{X}) \mid \text{Hom}(\mathbf{T}, \mathcal{F}) = 0\}.$$

Then $X_0 \cap X_1 = \{0\}$, X_0 and X_1 are both stable under extension, X_1 is stable under subobjects while X_0 is stable under quotients. Furthermore, from [Le], Cor. 4.7 we see that every indecomposable object is either in X_0 or in X_1 .

Recall the notations from Section 1.4. Let $\Delta_{\mathbf{l}}$ denote the set of roots of $\mathbf{l}$, and let $\hat{\Delta}_{\mathbf{f}}^{\geq 0} = \Delta_{\mathbf{l}} \oplus \mathbb{N}\delta$ be the set of roots of $\hat{\mathbf{f}}_{\geq 0}$.

Lemma 7.4. *The map $\mathcal{F} \mapsto [\mathcal{F}] \in K_0(\text{Coh}(\mathbb{X}_{\mathbf{p}, \Delta}))$ induces a bijection between the set of indecomposable vector bundles in X_0 and $\hat{\Delta}_{\mathbf{f}}^{\geq 0}$.*

Proof. Let $\mathcal{F}$ be an indecomposable vector bundle. Since $X_0 \cap X_1 = \{0\}$ and since every indecomposable vector bundle is either in X_0 or in X_1 , it is enough to show that $\mathcal{F} \notin X_1$ if and only if $[\mathcal{F}] \in \hat{\Delta}_{\mathbf{f}}^{\geq 0}$. Let us write $[\mathcal{F}] = \alpha' + d\delta + r[\mathcal{O}]$. It follows from Lemma 6.1 that

$$\langle \mathcal{O}, \mathcal{F} \rangle = r + d, \quad \langle \mathcal{O}(\vec{c}), \mathcal{F} \rangle = d.$$

In particular, if $d \geq 0$ then $\dim \text{Hom}(\mathbf{T}, \mathcal{F}) \geq \dim \text{Hom}(\mathcal{O}, \mathcal{F}) \geq \langle \mathcal{O}, \mathcal{F} \rangle > 0$, from which we deduce that $\mathcal{F} \in X_0$. On the other hand, if $d < 0$ then $\dim \text{Ext}^1(\mathbf{T}, \mathcal{F}) \geq \dim \text{Ext}^1(\mathcal{O}(\vec{c}), \mathcal{F}) \geq -\langle \mathcal{O}(\vec{c}), \mathcal{F} \rangle > 0$, from which we deduce that $\mathcal{F} \in X_1$. The Lemma is proved. $\diamond$

Proposition 7.3. *The algebra $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ contains the classes of all indecomposable vector bundles in X_0 .*

Proof. We argue again by induction on the rank. It is easy to see that a line bundle $\mathcal{O}(\vec{x})$ is in X_0 if and only if $\vec{x} \geq 0$. From Lemma 7.2 we deduce the statement of the Proposition for line bundles. Assume that $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ contains all indecomposable vector bundles of X_0 of rank at most r , and let $\mathcal{F} \in X_0$ be indecomposable of rank $r + 1$. Since $\text{Hom}(\mathbf{T}, \mathcal{F}) \neq 0$ there exists a line bundle $\mathcal{O}(\vec{x})$ with $\vec{x} \geq 0$, a vector bundle $\mathcal{G}$ and an exact sequence

$$0 \rightarrow \mathcal{O}(\vec{x}) \rightarrow \mathcal{F} \rightarrow \mathcal{G} \rightarrow 0.$$

Since $\mathcal{F} \in X_0$, the vector bundle $\mathcal{G}$ decomposes as a sum $\mathcal{G} = \mathcal{G}_1 \oplus \cdots \oplus \mathcal{G}_s$ of indecomposable vector bundles $\mathcal{G}_i \in X_0$ of rank at most r . Arguing as in Proposition 7.2, we see that $\mathcal{G} \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$. Now let us write

$$[\mathcal{O}(\vec{x})] \cdot [\mathcal{G}] = c_{\mathcal{F}}[\mathcal{F}] + R, \quad \text{for some } 0 \neq c_{\mathcal{F}} \in \mathbb{C}_v.$$

Since there is at most one indecomposable vector bundle with a given class in $K_0(\text{Coh}(\mathbb{X}))$ it follows that R is a sum of terms of the form $c_{\mathcal{H}}[\mathcal{H}]$ where H is a direct sum of indecomposable sheaves of rank at most r . Observe that these sheaves are in X_0 since X_0 is stable under extension. By the same reasoning as for $\mathcal{G}$ we conclude that $[\mathcal{H}] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$. Thus $[\mathcal{F}] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ and the induction is complete. $\diamond$

7.4. The following result is clear from the discussions of Section 4.2 and Section 4.3.

Lemma 7.5. *The Hall algebra $\mathbf{U}_{\mathbf{p}, \Delta}^{\infty}$ of the category of torsion sheaves on $\mathbb{X}$ is isomorphic to the subalgebra of $\mathbf{U}_v(\hat{\mathfrak{n}})$ generated by $\mathbf{H}_{p_s}$ for $s = 1, \dots, n$ and $H_{\star, r}$ for $r \geq 1$.*

In particular, for any $\alpha \in \hat{Q}$, we have the equality of dimensions

$$(7.6) \quad \dim(\mathbf{U}_{\mathbf{p}, \Delta}^{\infty}[\alpha]) = \dim(\mathbf{U}(\hat{\mathfrak{t}})[\alpha]).$$

7.5. We are now in position to prove (7.4). Let $\mathcal{F}_1, \dots, \mathcal{F}_s$ be any indecomposable vector bundles in X_0 , ordered so that $\mu(\mathcal{F}_1) \leq \dots \leq \mu(\mathcal{F}_s)$. Then

$$[\mathcal{F}_1] \cdot [\mathcal{F}_2] \cdots [\mathcal{F}_s] = c[\mathcal{F}_1 \oplus \cdots \oplus \mathcal{F}_s], \quad \text{for some } 0 \neq c \in \mathbb{C}_v.$$

Hence $[\mathcal{F}_1 \oplus \cdots \oplus \mathcal{F}_s] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$. In this way we obtain a linearly independent set. In particular, if $\dot{\mathbf{U}}_{\mathbf{p}, \Delta}^{\geq 0}$ is the subalgebra of $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ consisting of all classes of vector bundles then it follows from Lemma 7.4, Proposition 7.3 and the PBW theorem that, for any $\alpha \in \hat{Q}$, we have

$$\dim(\dot{\mathbf{U}}_{\mathbf{p}, \Delta}^{\geq 0}[\alpha]) = \dim(\mathbf{U}(\hat{\mathfrak{f}}_{\geq 0})[\alpha]).$$

On the other hand, since the category of torsion sheaves coincides with $\mathcal{C}_{\infty}$, it follows from Corollary 7.1 that the multiplication map

$$\dot{\mathbf{U}}_{\mathbf{p}, \Delta}^{\geq 0} \otimes \mathbf{U}_{\mathbf{p}, \Delta}^{\infty} \rightarrow \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$$

is injective. Thus, for any $\alpha \in \hat{Q}$, we have

$$\begin{aligned} \dim \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}[\alpha] &\geq \dim(\dot{\mathbf{U}}_{\mathbf{p}, \Delta}^{\geq 0} \otimes \mathbf{U}_{\mathbf{p}, \Delta}^{\infty})[\alpha] \\ &= \dim(\mathbf{U}(\hat{\mathfrak{f}}_{\geq 0}) \otimes \mathbf{U}(\hat{\mathfrak{t}})[\alpha]) \\ &= \dim(\mathbf{U}^{\geq 0}(\hat{\mathfrak{n}})[\alpha]), \end{aligned}$$

proving (7.4) as desired.

Remark. The set

$\mathbf{B} = \{[\mathcal{F}] \mid \mathcal{F} \text{ is a vector bundle or a torsion sheaf supported at exceptional points}\}$

can be considered as a natural analog of the PBW basis of the quantum Kac-Moody algebra $\mathbf{U}_q^+(\mathfrak{g})$ constructed using quivers (when $\mathfrak{g}$ is of finite type). However, in the present case, $\mathbf{B}$ is not a basis of $\mathbf{U}_v(\hat{\mathfrak{n}})$ but rather only of the (quantum analog of the) subalgebra generated by $\hat{\mathfrak{n}}_s^+$ for $s = 1, \dots, n$ and $e_{\star, t}$ for $t \in \mathbb{Z}$.

8. THE AFFINE CASE

In this section we prove Theorem 5.2 when Γ is an affine Dynkin diagram (necessarily of type $D_4^{(1)}$, $E_6^{(1)}$, $E_7^{(1)}$ or $E_8^{(1)}$), which we now assume is the case.

8.1. Let k be any field. The classification of indecomposable objects in $\text{Coh}(\mathbb{X})$ is obtained in [LM]. The main tool used there is the theory of mutations (as in e.g., [Bo]), which we now briefly recall. Let $\mathcal{C}$ be any hereditary k -linear category with finite-dimensional morphism and extension spaces. We assume that $\mathcal{C}$ has Serre duality, i.e that there exists an equivalence $\tau : \mathcal{C} \rightarrow \mathcal{C}$ and an isomorphism of bifunctors

$$(\text{Ext}^1(X, Y))^* \simeq \text{Hom}(Y, \tau X).$$

Let $\mathbf{A} = \{A_1, \dots, A_p\}$ be a family of pairwise nonisomorphic indecomposable objects in $\mathcal{C}$ such that $\text{End}(A_i) = k$, $\tau(A_i) \simeq A_{i+1 \bmod p}$ and such that $\text{Hom}(A_i, A_j) = 0$ if $i \neq j$. We denote by $\mathcal{A}$ the additive full subcategory of $\mathcal{C}$ generated by $\mathbf{A}$.

We define the functor $L_{\mathbf{A}} : \mathcal{C} \rightarrow \mathcal{C}$ of *left mutation* as follows. For any object F of $\mathcal{C}$ we set $\Sigma_{\mathbf{A}} F = \bigoplus_j \text{Hom}(A_j, F) \otimes_k A_j$ and we denote by $\sigma_F : \Sigma_{\mathbf{A}} F \rightarrow F$ the natural map. Then $L_{\mathbf{A}}(F)$ is defined by the exact sequence

$$\Sigma_{\mathbf{A}} F \rightarrow F \rightarrow L_{\mathbf{A}} F \rightarrow 0.$$

The *right mutation* $R_{\mathbf{A}} : \mathcal{C} \rightarrow \mathcal{C}$ is defined in a similar fashion. Let F be an object of $\mathcal{C}$ and set $\Omega_{\mathbf{A}} F = \bigoplus_j \text{Ext}^1(F, A_j)^* \otimes_k A_j$. The functors $\text{Hom}(\Omega_{\mathbf{A}} F, -)$ and $\text{Ext}^1(F, -)$ from $\mathcal{A}$ to $\text{mod } k$ are isomorphic. This gives rise to an exact sequence (the universal extension)

$$\mu_F : 0 \rightarrow \Omega_{\mathbf{A}} F \rightarrow R_{\mathbf{A}} F \rightarrow F \rightarrow 0$$

such that the connecting homomorphism $\text{Hom}(\Omega_{\mathbf{A}} F, A) \rightarrow \text{Ext}^1(F, A)$ is bijective for all $A \in \mathcal{A}$. Note that μ_F depends on a choice of isomorphism $\text{Hom}(\Omega_{\mathbf{A}} F, -) \simeq \text{Ext}^1(F, -)$, but $R_{\mathbf{A}} F$ doesn't. Moreover, the assignment $F \mapsto R_{\mathbf{A}} F$ is not functorial in general.

Finally, define full subcategories $\mathcal{A}^{\perp+}$ and $\mathcal{A}^{\perp-}$ of $\mathcal{C}$ by the following conditions :

$$\begin{aligned} \mathcal{A}^{\perp+} &= \{F \in \mathcal{C} \mid \text{Hom}(F, A_j) = 0 \text{ for all } j\}, \\ \mathcal{A}^{\perp-} &= \{F \in \mathcal{A}^{\perp+} \mid \sigma_F : \Sigma_{\mathbf{A}} F \rightarrow F \text{ is a monomorphism}\}. \end{aligned}$$

Theorem 8.1 ([LM]). *The functor of left mutations $L_{\mathbf{A}}$ induces an equivalence of categories $L : \mathcal{A}^{\perp-} \rightarrow \mathcal{A}^{\perp+}$, with inverse given by the right mutation $R_{\mathbf{A}}$. Moreover, $L_{\mathbf{A}}$ and $R_{\mathbf{A}}$ both commute with the equivalence τ .*

In particular, the restriction of $R_{\mathbf{A}}$ to $\mathcal{A}^{\perp+}$ is a functor.

Using the above methods, with $\mathcal{C} = \text{Coh}(\mathbb{X})$ and $\mathbf{A} = \{\mathcal{O}(i\vec{\omega})\}$, Lenzing and Meltzer proved the following result. Recall that $\mathcal{C}_q$ denotes the full subcategory consisting of the zero sheaf together with semistable sheaves of slope q . In particular, $\mathcal{C}_{\infty}$ is the subcategory of all torsion sheaves.

Theorem 8.2 ([LM]). *The following holds*

- i) *The categories $\mathcal{C}_q$ for $q \in \mathbb{Q} \cup \{\infty\}$ are all canonically equivalent.*
- ii) *For $q > 0$, left and right mutations with respect to $\mathbf{A} = \{\mathcal{O}(i\vec{\omega})\}$ induce inverse equivalences*

$$\mathcal{C}_q \xrightarrow{R_{\mathbf{A}}} \mathcal{C}_{\frac{q}{q+1}} \xrightarrow{L_{\mathbf{A}}} \mathcal{C}_q.$$

In particular, the categorie $\mathcal{C}_q$ are described, like $\mathcal{C}_{\infty}$, in terms of categories of representations of cyclic quivers (see Section 2.4).

8.2. Recall that the symmetric bilinear form $(\cdot, \cdot)$ on $K_0(\text{Coh}(\mathbb{X}))$ coincides with the Cartan form on $\widehat{Q}$ (Proposition 5.1). This, together with Corollary 1.1 and [LM], Theorem 4.6. implies the following result. We denote by $\widehat{\Delta}_{\mathbf{n}}$ the set of roots of $\widehat{\mathbf{n}}$.

Proposition 8.1. *There is an indecomposable sheaf $\mathcal{F} \in \text{Coh}(\mathbb{X})$ such that $[\mathcal{F}] = \alpha \in \widehat{Q}$ if and only if $\alpha \in \widehat{\Delta}_{\mathbf{n}}$. Such a sheaf is unique (up to isomorphism) if and only if α is a real root. Moreover, $\mathcal{F}$ is a torsion sheaf (resp. a locally free sheaf) if and only if $[\mathcal{F}]$ is a root of $\widehat{\mathfrak{t}}$ (resp. a root of $\widehat{\mathfrak{f}}$).*

Combining Theorem 8.2 and Proposition 8.1 we obtain the following description of the structure of the set $\widehat{\Delta}_{\mathbf{n}}$. Consider linear forms $\mathbf{d} : \widehat{Q} \rightarrow \mathbb{Z}$ and $\mathbf{r} : \widehat{Q} \rightarrow \mathbb{Z}$ corresponding to the degree and rank forms on $K_0(\text{Coh}(\mathbb{X}))$. For $\alpha \in \widehat{Q}$ set $\mu(\alpha) = \frac{\mathbf{d}(\alpha)}{\mathbf{r}(\alpha)} \in \mathbb{Q} \cup \{\infty\}$ and for $q \in \mathbb{Q} \cup \{\infty\}$ put $\widehat{\Delta}_q = \{\alpha \in \widehat{\Delta}_{\mathbf{n}} \mid \mu(\alpha) = q\}$ and $\widehat{\mathbf{n}}_q = \bigoplus_{\alpha \in \widehat{\Delta}_q} \widehat{\mathbf{n}}[\alpha]$.

Corollary 8.1. *The following holds :*

- i) Each $\widehat{\mathbf{n}}_q$ is a Lie subalgebra, $\widehat{\mathbf{n}} = \bigoplus_q \widehat{\mathbf{n}}_q$ (as graded vector spaces) and $\widehat{\mathbf{n}}_{\infty} = \widehat{\mathfrak{t}}$,
- ii) For each q there is a canonical isomorphism $u_q : \widehat{\Delta}_q \simeq \widehat{\Delta}_{\infty}$ which lifts to a vector space isomorphism $\widehat{\mathbf{n}}_q \simeq \widehat{\mathbf{n}}_{\infty}$.
- iii) The automorphism $\mathcal{F} \mapsto \mathcal{F}(\vec{\omega})$ of $\text{Coh}(\mathbb{X})$ induces, for each q an automorphism τ of $\widehat{\Delta}_q$ of order p .

Proof. Assertions i) and iii) are immediate, as is the first claim of ii). Observe that u_q is induced by equivalences of categories, and thus compatible with the Euler form. In particular, u_q maps real roots to real roots and imaginary roots to imaginary roots. But all real roots of $\widehat{\mathbf{n}}$ have multiplicity one and all imaginary roots have multiplicity $|\Gamma|$. Therefore, u_q lifts (in a noncanonical way) to a vector space isomorphism $\widehat{\mathbf{n}}_q \simeq \widehat{\mathbf{n}}_{\infty}$. $\diamond$

Remarks. It is possible to show that in fact $\widehat{\mathbf{n}}_q \simeq \widehat{\mathbf{n}}_{\infty}$ as Lie algebras. We won't need this result.

We will call a root $\alpha \in \widehat{\Delta}_q$ *simple* if it corresponds, under the equivalence $\widehat{\Delta}_q \simeq \widehat{\Delta}_{\infty}$, to the class of a simple torsion sheaf $[S_i^{(s)}]$ (supported at an exceptional point). Note that all simple roots are real. In addition, the simple roots $\alpha \in \widehat{\Delta}_q$ decompose as a union of n τ -orbits of size $p_1, \dots, p_n$ respectively, and the set of simple sheaves in $\mathcal{C}_q$ is naturally in bijection with $\mathbb{Z}/p_1\mathbb{Z} \sqcup \dots \sqcup \mathbb{Z}/p_n\mathbb{Z}$. On the other hand, there exists a unique minimal imaginary root $\delta_q \in \widehat{\Delta}_q$ of which all other imaginary roots are multiples.

8.3. We now introduce some exhaustive filtrations on $\mathbf{U}(\widehat{\mathbf{n}})$, $\mathbf{U}_v(\widehat{\mathbf{n}})$ and $\mathbf{U}_{\mathbf{p}, \Delta}^{(k)}$ respectively. Let $\widehat{\mathbf{n}}_{\geq 0}$ be the subalgebra of $\widehat{\mathbf{n}}$ generated by $\widehat{\mathfrak{t}}$ and by the root vectors corresponding to the simple roots $\alpha_*, \tau(\alpha_*), \dots, \tau^{p-1}(\alpha_*)$ in $\widehat{\Delta}_0$. We set $\mathbf{U}^{\geq 0}(\widehat{\mathbf{n}}) = \mathbf{U}(\widehat{\mathbf{n}}_{\geq 0})$.

Lemma 8.1. *The algebra $\mathbf{U}^{\geq 0}(\widehat{\mathbf{n}})$ has finite-dimensional weight spaces.*

Proof. For $m \in \mathbb{Z}$, let us denote by $\widehat{\mathbf{u}}_{\geq m}$ the subalgebra of $\widehat{\mathbf{n}}$ generated by $\widehat{\mathfrak{t}}$ and $e_{*, t}$ for $t \geq m$. This defines an exhaustive filtration of $\widehat{\mathbf{n}}$, and the proof of Lemma 7.1 shows that $\mathbf{U}(\widehat{\mathbf{u}}_{\geq m})$ has finite-dimensional weight spaces for any m . Now choose $m \ll 0$ such that $\widehat{\mathbf{u}}_{\geq m}$ contains the root vectors of weight $\alpha_*, \tau(\alpha_*), \dots, \tau^{p-1}(\alpha_*)$. Then, for any $\alpha \in \widehat{Q}$ we have

$$\dim \mathbf{U}^{\geq 0}(\widehat{\mathbf{n}})[\alpha] \leq \dim \mathbf{U}(\widehat{\mathbf{u}}_{\geq m})[\alpha] < \infty.$$

◇

The following result is proved in the same way as Lemma 7.2.

Lemma 8.2. *The algebra $\mathbf{U}_{\mathbf{p},\lambda}^{(k)}$ contains the classes of all line bundles $\mathcal{O}(\vec{x})$.*

Let $E(\vec{x}) \in \mathbf{U}_v(\widehat{\mathbf{n}})$ be the element satisfying $\Phi(E(\vec{x})) = [\mathcal{O}(x)]$ provided by the proof of Lemma 7.2. Let $\mathbf{U}_{\mathbf{p},\lambda}^{\geq 0,(k)}$ (resp. $\mathbf{U}_v^{\geq 0}(\widehat{\mathbf{n}})$) be generated by the classes of the simple torsion sheaves $[S_i^{(s)}]$, by T_r for $r \geq 1$ and by the classes of the line bundles $[\mathcal{O}(i\vec{\omega})]$ for $i = 0, \dots, p-1$ (resp. by $\mathbf{H}_{p_i}$ for $i = 1, \dots, n$ and by $E(i\vec{\omega})$ for $i = 0, \dots, p-1$). It is clear that $\Phi(\mathbf{U}_v^{\geq 0}(\widehat{\mathbf{n}})) = \mathbf{U}_{\mathbf{p},\lambda}^{\geq 0,(k)}$.

Let us denote by σ the automorphism of $\mathbf{U}(\widehat{\mathbf{n}})$ (resp. of $\mathbf{U}_v(\widehat{\mathbf{n}})$) defined by $\sigma(e_{\star,t}) = e_{\star,t+1}$ and the identity on all other generators (resp. by $\sigma(E_{\star,t}) = E_{\star,t+1}$ and the identity on all other generators). Let us also denote by the same letter the automorphism of $\mathbf{U}_{\mathbf{p},\lambda}^{(k)}$ defined by $\sigma([\mathcal{F}]) = [\mathcal{F}(\vec{c})]$ for any $\mathcal{F} \in \text{Coh}(\mathbb{X})$. It is clear that σ commutes with the morphism $\Phi^{(k)} : \mathbf{U}_v(\widehat{\mathbf{n}}) \rightarrow \mathbf{U}_{\mathbf{p},\lambda}^{(k)}$. Finally, for $m \in \mathbb{Z}$ we put

$$\mathbf{U}^{\geq m}(\widehat{\mathbf{n}}) = \sigma^m(\mathbf{U}^{\geq 0}(\widehat{\mathbf{n}})), \quad \mathbf{U}_v^{\geq m}(\widehat{\mathbf{n}}) = \sigma^m(\mathbf{U}_v^{\geq 0}(\widehat{\mathbf{n}})), \quad \mathbf{U}_{\mathbf{p},\lambda}^{\geq m,(k)} = \sigma^m(\mathbf{U}_{\mathbf{p},\lambda}^{\geq 0,(k)}).$$

Arguing in the same way as in Section 7.1, we see that Theorem 5.2 will follow from the set of inequalities

$$(8.1) \quad \dim_{\mathbb{C}} \mathbf{U}_{\mathbf{p},\lambda}^{\geq 0,(k)}[\alpha] \geq \dim_{\mathbb{C}} \mathbf{U}^{\geq 0}(\widehat{\mathbf{n}})[\alpha]$$

for all finite fields k and all $\alpha \in \widehat{Q}$. This set of inequalities is proved in the rest of this Section.

8.4. From now on we fix the finite field k and drop the superscript (k) in all notations. The aim of this paragraph is to prove the following result.

Proposition 8.2. *$\mathbf{U}_{\mathbf{p},\lambda}^{\geq 0}$ contains the classes of all simple sheaves in $\mathcal{C}_q$ for $q > 0$.*

We start with some preliminary technical results. Define a sequence F_m for $m \geq 1$ (the *Farey sequence*) inductively as follows. We set $F_1 = \{\frac{0}{1}, \frac{1}{0}\}$ and if $F_m = \{\frac{0}{1}, \frac{a_1}{b_1}, \dots, \frac{a_r}{b_r}, \frac{1}{0}\}$ then

$$F_{m+1} = \left\{ \frac{0}{1}, \frac{a_1}{b_1+1}, \frac{a_1}{b_1}, \dots, \frac{a_i}{b_i}, \frac{a_i+a_{i+1}}{b_i+b_{i+1}}, \frac{a_{i+1}}{b_{i+1}}, \dots, \frac{1}{0} \right\}.$$

In the above, we treat the fraction $\frac{1}{0}$ as representing ∞ . The following facts are well-known about the sequences F_m :

- i) Each new term $\frac{a_i+a_{i+1}}{b_i+b_{i+1}}$ appearing in F_{m+1} is a reduced fraction.
- ii) If $\frac{a_i}{b_i}$ and $\frac{a_{i+1}}{b_{i+1}}$ are consecutive entries in some F_m then $b_i a_{i+1} - b_{i+1} a_i = 1$.
- iii) Every positive rational number belongs to F_m for $m \gg 0$.

Proposition 8.3. *Let $\frac{a}{b}$ and $\frac{c}{d}$ be consecutive entries in some F_m . Let $\mathbf{U} = \{U_i\}_{i \in I}$ be simple sheaves in $\mathcal{C}_{\frac{a}{b}}$ forming a single τ -orbit of size p . Let $\mathcal{U}$ be the additive closure of $\mathbf{U}$. Then*

- i) For any $i \in I$, U_i is a vector bundle of rank b and of degree a ,
- ii) $\mathcal{C}_{\frac{a+c}{b+d}} \subset \mathcal{U}^\perp$ and the right mutation $R_{\mathbf{U}}$ restricts to an equivalence $\mathcal{C}_{\frac{c}{d}} \xrightarrow{\sim} \mathcal{C}_{\frac{a+c}{b+d}}$.

Let us denote by $\{V_j\}_{j \in J}$ the collection of all simple sheaves in $\mathcal{C}_{\frac{c}{d}}$. Then for any $j \in J$,

$$\text{iii) } \sum_i \dim \text{Ext}^1(V_j, U_i) = \frac{p}{c_j}, \quad \dim \text{Ext}^1(V_j, U_i) \leq 1 \quad \text{for all } i,$$

where $c_j \in \{p_1, \dots, p_n\}$ is the size of the τ -orbit of V_j .

Proof. We will give a proof by induction. The result is easily checked for F_1 (see [LM], Proposition 3.9 for ii)). Let us assume that properties i) and iii) are proved for all consecutive entries of F_l . We will prove ii) for entries in F_l and i) and iii) for entries in F_{l+1} .

Let $\frac{a}{b}$ and $\frac{c}{d}$ be consecutive entries in F_l , and let $\mathbf{U} = \{U_i\}_{i \in I}$ and $\{V_j\}_{j \in J}$ be as above.

Claim 8.1. *We have $\mathcal{C}_{\frac{a+c}{b+d}} \subset \mathcal{U}^\perp$.*

Proof. Let $\mathcal{F}$ be an indecomposable sheaf in $\mathcal{C}_{\frac{a+c}{b+d}}$. Since $\frac{a}{b} < \frac{a+c}{b+d}$ we have $\mathcal{F} \in \mathcal{U}^\perp$. Consider the morphism $\sigma_{\mathcal{F}} : \Sigma_{\mathbf{U}} \mathcal{F} \rightarrow \mathcal{F}$ and let $\mathcal{G}$ be the image of $\sigma_{\mathcal{F}}$. Thus $\sigma_{\mathcal{G}} : \Sigma_{\mathbf{U}} \mathcal{G} = \Sigma_{\mathbf{U}} \mathcal{F} \rightarrow \mathcal{G}$ is an epimorphism. We may assume that $\mathcal{G}$ is indecomposable. We claim that $\text{Hom}(\mathcal{G}, U_i) \neq 0$ for some $i \in I$. Indeed, if not then by Serre duality $\text{Ext}^1(U_i, \mathcal{G}) = 0$ for all i and using the Riemann-Roch theorem we obtain

$$\begin{aligned} \sum_i \dim \text{Hom}(U_i, \mathcal{G}) &= \sum_i \langle U_i, \mathcal{G} \rangle \\ (8.2) \quad &= \mathbf{r}(U_i) \mathbf{d}(\mathcal{G}) - \mathbf{d}(U_i) \mathbf{r}(\mathcal{G}) \\ &= b \mathbf{d}(\mathcal{G}) - a \mathbf{r}(\mathcal{G}). \end{aligned}$$

On the other hand, we have

$$(8.3) \quad \sum_i \dim \text{Hom}(U_i, \mathcal{G}) = \frac{\mathbf{r}(\Sigma_{\mathbf{U}} \mathcal{G})}{b} \leq \frac{\mathbf{r}(\mathcal{G})}{b}.$$

Combining (8.2) and (8.3) we deduce that $\mu(\mathcal{G}) \geq \frac{a}{b} + \frac{1}{b^2}$. But since $\mathcal{G} \hookrightarrow \mathcal{F}$ we have

$$\mu(\mathcal{G}) \leq \mu(\mathcal{F}) = \frac{a+c}{b+d} = \frac{a}{b} + \frac{bc-ad}{bd} = \frac{a}{b} + \frac{1}{b(b+d)},$$

a contradiction. Thus there exists a nonzero map $\gamma : \mathcal{G} \rightarrow U_i$. But then γ is automatically an isomorphism, and $\sigma_{\mathcal{G}}$ is also an isomorphism. This completes the proof of Claim 8.1. $\diamond$

Now, since V_j is of rank d and of degree c , the Riemann-Roch theorem implies that

$$(8.4) \quad \mathbf{d}(R_{\mathbf{U}} V_j) = c + (bc - ad)a = a + c,$$

$$(8.5) \quad \mathbf{r}(R_{\mathbf{U}} V_j) = d + (bc - ad)b_i = b + d.$$

In particular, $\mu(R_{\mathbf{U}} V_j) = \frac{a+c}{b+d}$. Hence from the above claim we deduce that $\{R_{\mathbf{U}} V_j\}_{j \in J}$ forms a complete set of simple sheaves in $\mathcal{C}_{\frac{a+c}{b+d}}$ and ii) follows. Note that i) for $\frac{a+c}{b+d}$ is now a consequence of (8.4) and (8.5).

It remains to prove iii) for the pairs $\{\frac{a}{b}, \frac{a+c}{b+d}\}$ and $\{\frac{a+c}{b+d}, \frac{c}{d}\}$. This is taken care of by the following claims.

Claim 8.2. *For all $j \in J$ we have*

$$(8.6) \quad \sum_i \dim \text{Ext}^1(R_{\mathbf{U}} V_j, U_i) = \frac{p}{c_j}, \quad \dim \text{Ext}^1(R_{\mathbf{U}} V_j, U_i) \leq 1 \quad \text{for all } i.$$

Proof. By definition, we have

$$[R_{\mathbf{U}} V_j] = [V_j] + \sum_k \dim \text{Ext}^1(V_j, U_k) [U_k] = [V_j] - \sum_k \langle V_j, U_k \rangle [U_k]$$

since $V_j \in \mathcal{C}_{q'}$, $U_k \in \mathcal{C}_q$ and $q < q'$, so that $\text{Hom}(V_j, U_k) = 0$. Observe that $R_{\mathbf{U}}V_j$ is indecomposable by Theorem 8.1. Hence $R_{\mathbf{U}}V_j \in \mathcal{C}_{q''}$ for some $q < q'' < q'$. Therefore,

$$\dim \text{Ext}^1(R_{\mathbf{U}}V_j, U_i) = -\langle R_{\mathbf{U}}V_j, U_i \rangle = -\langle V_j, U_i \rangle + \sum_k \langle V_j, U_k \rangle \langle U_k, U_i \rangle.$$

But by Theorem 8.2 and Lemma 6.1, we have

$$(8.7) \quad \langle U_k, U_i \rangle = \begin{cases} 1 & \text{if } U_k = U_i \\ -1 & \text{if } \tau(U_k) = U_i \\ 0 & \text{otherwise} \end{cases}.$$

Hence $\dim \text{Ext}^1(R_{\mathbf{U}}V_j, U_i) = \langle V_j, \tau^{-1}U_i \rangle$, and (8.6) is proved. $\diamond$

Similarly, let $K \subset J$ be such that $\{V_k \mid k \in K\}$ forms a single τ -orbit of size p .

Claim 8.3. *For all $j \in J$ we have*

$$(8.8) \quad \sum_k \dim \text{Ext}^1(V_j, R_{\mathbf{U}}V_k) = \frac{p}{c_j}, \quad \dim \text{Ext}^1(V_j, R_{\mathbf{U}}V_k) \leq 1 \quad \text{for all } k.$$

Proof. As in the previous lemma,

$$\dim \text{Ext}^1(V_j, R_{\mathbf{U}}V_k) = -\langle V_j, R_{\mathbf{U}}V_k \rangle = -\langle V_j, R_{\mathbf{U}}V_k \rangle + \sum_i \langle V_j, U_i \rangle \langle V_k, U_i \rangle.$$

Since the τ -orbit of V_k is of size p , the induction hypothesis on $\text{Ext}^1(V_k, U_i)$ implies that there exists $i(k) \in I$ such that

$$\langle V_k, U_{i(k)} \rangle = 1, \quad \langle V_k, U_{i'} \rangle = 0 \text{ for all } i' \neq i(k).$$

As a consequence,

$$\dim \text{Ext}^1(V_j, R_{\mathbf{U}}V_k) = -\langle V_j, R_{\mathbf{U}}V_k \rangle + \langle V_j, U_{i(k)} \rangle$$

Finally, it is easy to see that the assignment $k \mapsto i(k)$ is a bijection between K and I . Using (8.7) we deduce equation (8.8).

This concludes the proof of Claim 8.3 and of Proposition 8.3. $\diamond$

We are now in position to prove Proposition 8.2. We argue again by induction. By definition, $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ contains the classes of all the simple sheaves in $\mathcal{C}_{\infty}$ and the classes of simple sheaves $\mathcal{O}(i\vec{\omega})$. Assume that $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ contains the classes of all simple sheaves of $\mathcal{C}_q$ if $q > 0$ and q appears in F_l . Let $\frac{a}{b}$ and $\frac{c}{d}$ be two consecutive entries in F_l . Pick a collection $\mathbf{U} = \{U_i\}_{i \in I} \subset \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ of simple sheaves in $\mathcal{C}_{\frac{c}{d}}$ forming a single τ -orbit of size p . Let V be any simple sheaf of $\mathcal{C}_{\frac{a}{b}}$. From Proposition 8.2, iii), we deduce that there exists $I_0 \subset I$ such that $\Sigma_{\mathbf{U}}V = \bigoplus_{i \in I_0} U_i$. Recall that $R_{\mathbf{U}}V$ is the universal extension of V with respect to $\mathcal{U}$. Moreover, it follows from Proposition 8.3, iii) and Serre duality that

$$\dim \text{Hom}(U_i, R_{\mathbf{U}}V) = \langle U_i, R_{\mathbf{U}}V \rangle = -\langle R_{\mathbf{U}}V, \tau U_i \rangle \in \{0, 1\}.$$

We conclude that for any subset $J \subset I_0$ there exists a unique indecomposable object O_J fitting in an exact sequence

$$0 \rightarrow \bigoplus_{i \in J} U_i \rightarrow O_J \rightarrow V \rightarrow 0$$

(specifically, $O_J \simeq R_{\mathbf{U}}V / \bigoplus_{j \notin J} U_j$).

Claim 8.4. $[O_J] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ for any $J \subset I_0$.

Proof. We argue by induction. Assume that $[O_J] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ for any subset J of size at most k , and take $J \subset I_0$ with $|J| = k$. Observe that for any proper subset $J \subset I$ the class $[\bigoplus_{j \in J} U_j]$ is in $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ (this can easily be deduced from the fact that the Serre subcategory of $\text{Coh}(\mathbb{X})$ generated by $\mathbf{U}$ is equivalent to the category of representations of the cyclic quiver $A_{p-1}^{(1)}$). Thus,

$$(8.9) \quad \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0} \ni [\bigoplus_{j \in J} U_j] \cdot [V] = c_J [O_J] + \sum_{K \subset J, K \neq J} c_K [O_K \oplus \bigoplus_{l \notin K} U_l]$$

for some nonzero $c_J, c_K \in \mathbb{C}_v$. By the induction hypothesis, $[O_K] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ for all K , and thus $[O_K \oplus \bigoplus_{l \notin K} U_l] = c [O_K] \cdot [\bigoplus_{l \notin K} U_l] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$. We conclude that $[O_J] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$, proving the claim. $\diamond$

In particular, taking $J = I_0$ we see that $[R_{\mathbf{U}} V] \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$. Finally, by Proposition 8.2 ii), $R_{\mathbf{U}} V$ is a simple sheaf in $\mathcal{C}_{\frac{a+c}{b+d}}$ and every simple sheaf in $\mathcal{C}_{\frac{a+c}{b+d}}$ arises in this way. Thus $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ contains the classes of all simple sheaves in $\mathcal{C}_q$ for $q > 0$ arising in F_{l+1} . This completes the induction and the proof of Proposition 8.3. $\diamond$

8.4. In this paragraph, we construct certain elements in $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ corresponding to imaginary root vectors of $\hat{\mathfrak{n}}$. We first introduce some useful notation. For any $q \in \mathbb{Q}$ we denote by $\mathcal{C}_{>q}$ (resp. $\mathcal{C}_{<q}$) the additive subcategory of $\text{Coh}(\mathbb{X})$ generated by $\mathcal{C}_{q'}$ for $q' > q$ (resp. $q' < q$). Also, if $\mathcal{F}$ is any coherent sheaf and $u \in \mathbf{H}_{\text{Coh}(\mathbb{X})}$ then we let $c_{\mathcal{F}}(u)$ be the coefficient of $[\mathcal{F}]$ in u . Finally, we will say that a sheaf $\mathcal{F} \in \mathcal{C}_q$ is a λ -sheaf if $\mathcal{F} = \mathcal{X}_1 \oplus \dots \oplus \mathcal{X}_l$ where $\mathcal{X}_1, \dots, \mathcal{X}_l$ are indecomposable sheaves in $\mathcal{C}_q$ corresponding, under the equivalence $\mathcal{C}_q \simeq \mathcal{C}_{\infty}$ to simple torsion sheaves of degree $\lambda_1 \delta, \dots, \lambda_l \delta$ respectively.

Define elements $T_r^{(q)} \in \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ for $q \in \mathbb{Q}$, $q > 0$ and $r \geq 1$ as follows. Set $T_r^{\infty} = T_r$. Assume that $T_r^{(q)}$ has been defined for all q belonging to the sequence F_l . Let $\frac{a}{b}$ and $\frac{c}{d}$ be consecutive entries in F_l , and let $\mathbf{U} = \{U_i\}_{i \in \mathbb{Z}/p\mathbb{Z}}$ be any collection of simple sheaves in $\mathcal{C}_{\frac{a}{b}}$ forming a single τ -orbit. We put

$$T_r^{(\frac{a+c}{b+d})} = [U_1]^r \cdot [U_2]^r \cdots [U_p]^r \cdot T_r^{(\frac{c}{d})}.$$

This definition depends on the choice (and ordering) of the U_i but this will not be important for us. For any $q > 0$ and any partition $\lambda = (\lambda_1 \geq \dots \geq \lambda_l)$ we define $T_{\lambda}^{(q)} = T_{\lambda_1}^{(q)} \cdots T_{\lambda_l}^{(q)}$.

Claim 8.5. *We have $c_{\mathcal{F}}(T_{\lambda}^{(q)}) \neq 0$ for any λ -sheaf $\mathcal{F}$ in $\mathcal{C}_q$.*

Proof. Let us first consider the case when $\lambda = \lambda_1$, and argue by induction. The statement is clear for $q \in F_1$. Assume that it holds for all $q \in F_l$ and let $\frac{a}{b}, \frac{c}{d}$ and $\{U_i\}_{i \in \mathbb{Z}/p\mathbb{Z}}$ have the same meaning as above. Let $\mathcal{F} \in \mathcal{C}_{\frac{a}{b}}$ be a λ -sheaf. From Proposition 8.3, iii) we see that $\Omega_{\mathbf{U}}(\mathcal{F}) = U_1^{\oplus r} \oplus \dots \oplus U_p^{\oplus r}$. It remains to notice that

$$c_{[U_1^{\oplus r} \oplus \dots \oplus U_p^{\oplus r}]}([U_1]^r \cdot [U_2]^r \cdots [U_p]^r) \neq 0.$$

This proves the claim for λ of length one. The general case easily follows. $\diamond$

By Section 2.4 and Section 4.2, the subalgebra $\mathbf{C}_q$ of $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ generated by the simple sheaves in $\mathcal{C}_q$ is isomorphic to $\mathbf{U}_v^+(\widehat{\mathfrak{sl}}_{p_1}) \otimes \dots \otimes \mathbf{U}_v^+(\widehat{\mathfrak{sl}}_{p_n})$. Let $\mathcal{B}$ be a basis of the latter algebra consisting of weight vectors, and let $\mathcal{B}^{(q)}$ be the corresponding basis of $\mathbf{C}_q$.

Lemma 8.3. *For any $q > 0$ and any sheaf $\mathcal{F}$ appearing in $T_r^{(q)}$ we have $[\mathcal{F}] = r\delta_q \in K_0(\text{Coh}(\mathbb{X})) \simeq \widehat{Q}$.*

Proof. We argue again by induction. Assume the statement in the Lemma for all q appearing in F_l and let $\frac{a}{b}, \frac{c}{d}$ be consecutive entries in F_l and let $\mathbf{U} = \{U_i\}_{i \in \mathbb{Z}/p\mathbb{Z}}$ be any collection of simple sheaves in $\mathcal{C}_{\frac{a}{b}}$ forming a single τ -orbit. It is enough to observe that

$$\delta_{\frac{a+c}{b+d}} = [U_1] + \cdots + [U_p] + \delta_{\frac{c}{d}}$$

since the right mutation $R_{\mathbf{U}}$ defines an equivalence $\mathcal{C}_{\frac{c}{d}} \simeq \mathcal{C}_{\frac{a+c}{b+d}}$. $\diamond$

Proposition 8.4. *The elements $\{b^{(q)} \cdot T_{\lambda}^{(q)} \mid b^{(q)} \in \mathcal{B}^{(q)}, \lambda \in \Pi\}$ are linearly independent.*

Proof. Let $\text{Coh}^{ss}(\mathbb{X})$ denote the set of semistable coherent sheaves on $\mathbb{X}$. Set

$$I'_q = \bigoplus_{\substack{\mu(\mathcal{F})=q \\ \mathcal{F} \notin \text{Coh}^{ss}(\mathbb{X})}} \mathbb{C}[\mathcal{F}] \subset \mathbf{H}_{\text{Coh}(\mathbb{X})},$$

$$I_q^{ss} = \bigoplus_{\substack{\mu(\mathcal{F})=q \\ \mathcal{F} \in \text{Coh}^{ss}(\mathbb{X})}} \mathbb{C}[\mathcal{F}] \subset \mathbf{H}_{\text{Coh}(\mathbb{X})}.$$

If $[\mathcal{F}] \in I'_q$ then $\mathcal{F} = \mathcal{F}^- \oplus \mathcal{F}^0 \oplus \mathcal{F}^+$ where $\mathcal{F}^- \in \mathcal{C}_{<q}$ and $\mathcal{F}^+ \in \mathcal{C}_{>q}$ are both nonzero and where $\mathcal{F}^0 \in \mathcal{C}_q$.

Lemma 8.4. *$I'_q \oplus I_q^{ss}$ is a subalgebra of $\mathbf{H}_{\text{Coh}(\mathbb{X})}$, and I'_q is an ideal in $I'_q \oplus I_q^{ss}$.*

Proof. The first statement is obvious. For the second, observe that if $[\mathcal{G}] \in I_q^{ss}$ and $[\mathcal{F}] = [\mathcal{F}^- \oplus \mathcal{F}^0 \oplus \mathcal{F}^+] \in I'_q$ then by the HN filtration there can be no monomorphism $\mathcal{F}^- \oplus \mathcal{F}^0 \oplus \mathcal{F}^+ \hookrightarrow \mathcal{G}$ or epimorphism $\mathcal{G} \twoheadrightarrow \mathcal{F}^- \oplus \mathcal{F}^0 \oplus \mathcal{F}^+$. $\diamond$

We will write π_{ss} for the projection. $I_q^{ss} \oplus I'_q \rightarrow I_q^{ss}$. Note that $T_r^{(q)} \in I_q^{ss} \oplus I'_q$. By Lemma 8.4 we have, for any partition $\lambda = (\lambda_1 \geq \cdots \geq \lambda_l)$,

$$(8.10) \quad \pi_{ss}(T_{\lambda}^{(q)}) = \pi_{ss}(T_{\lambda_1}^{(q)}) \cdots \pi_{ss}(T_{\lambda_l}^{(q)}).$$

From this and Claim 8.5 we deduce the following result.

Claim 8.6. *Let $\mu > \lambda$ and let $\mathcal{F}$ be any μ -sheaf. Then $c_{\mathcal{F}}(T_{\lambda}^{(q)}) = 0$.*

Now, to prove that the set $\{b^{(q)} \cdot T_{\lambda}^{(q)}\}$ is linearly independent, it is enough to consider their image under the projection π_{ss} . Let us call a sheaf $\mathcal{G} \in \mathcal{C}_q$ *exceptional* if it corresponds, under the equivalence $\mathcal{C}_q \simeq \mathcal{C}_{\infty}$, to a torsion sheaf supported at exceptional points. In particular, if $\mathcal{B}^{(q)} \ni b^{(q)} = \sum_i c_i [\mathcal{G}_i]$ then all $\mathcal{G}_i$ are exceptional. Observe that if $\mathcal{G}$ is exceptional and if $\mathcal{F}$ is a λ -sheaf then (up to a scalar) $[\mathcal{G}] \cdot [\mathcal{F}] = c[\mathcal{F} \oplus \mathcal{G}]$. Proposition 8.4 now easily follows from Claim 8.6 and from the fact that the elements of $\mathcal{B}^{(q)}$ are linearly independent. $\diamond$

8.6. Let $\mathcal{I}$ denote the collection of all tuples

$$\sigma = \{q_1, \dots, q_l; d_1, \dots, d_l \mid q_i, d_i \in \mathbb{Q}^+, q_1 > q_2 > \cdots > q_l\}.$$

We introduce a total order on $\mathcal{I}$ as follows. $\sigma = \{q_i; d_i\} \succ \sigma' = \{q'_i; d'_i\}$ if there exists $j \geq 0$ such that $q_1 = q'_1, d_1 = d'_1, \dots, q_j = q'_j, d_j = d'_j$ and $q_{j+1} > q'_{j+1}$ or $q_{j+1} = q'_{j+1}$ and $d_i > d'_{i+1}$.

Let us associate to any coherent sheaf $\mathcal{F}$ the element $\sigma(\mathcal{F}) \in \mathcal{I}$ defined as

$$\sigma(\mathcal{F}) = \{q_1, \dots, q_l; \mathbf{d}(\mathcal{F}_1), \dots, \mathbf{d}(\mathcal{F}_l)\}$$

where $\mathcal{F} = \mathcal{F}_1 \oplus \cdots \oplus \mathcal{F}_l$ with $\mathcal{F}_i \in \mathcal{C}_{q_i}$ and $q_1 > \cdots > q_l$.

Proposition 8.5. Fix $q_1 > \dots > q_l > 0$ and for $1 \leq i \leq l$ let $b^{(q_i)} \in \mathcal{B}^{(q_i)}$ and $\lambda_i \in \Pi$. We have

$$(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}) \dots (b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \in \pi_{ss}(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}) \dots \pi_{ss}(b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \oplus \bigoplus_{\sigma(\mathcal{F}) \succ \gamma} \mathbb{C}[\mathcal{F}],$$

where $\gamma = \{q_1, \dots, q_l; \mathbf{d}(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}), \dots, \mathbf{d}(b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)})\}$.

Proof. We give a proof by induction. The statement is clear for $l = 1$. Assume that the Proposition is true for a product of $l - 1$ terms, and let us set

$$A = (b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}) \dots (b^{(q_{l-1})} \cdot T_{\lambda_{l-1}}^{(q_{l-1})}).$$

The next statement is a simple consequence of the HN filtration and of the definition of σ :

Lemma 8.5. Let $\mathcal{F}$ and $\mathcal{G}$ be any coherent sheaves in $\mathcal{C}_{>0}$. Then

$$[\mathcal{F}] \cdot [\mathcal{G}] \in \bigoplus_{\sigma(\mathcal{H}) \succ \sigma(\mathcal{F})} \mathbb{C}[\mathcal{H}].$$

In particular, we deduce that

$$A \cdot (b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \in \pi_{ss}(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}) \dots \pi_{ss}(b^{(q_{l-1})} \cdot T_{\lambda_{l-1}}^{(q_{l-1})}) \cdot (b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \oplus \bigoplus_{\sigma(\mathcal{F}) \succ \gamma} \mathbb{C}[\mathcal{F}].$$

But

$$b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)} \in \pi_{ss}(b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \oplus \bigoplus_{\sigma(\mathcal{H}) \succ \gamma'} \mathbb{C}[\mathcal{H}],$$

where $\gamma' = \{q_l; \mathbf{d}(b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)})\}$. The Proposition easily follows. $\diamond$

Corollary 8.2. The elements of $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$

$$\{(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}) \dots (b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \mid q_1 > \dots > q_l, b^{(q_i)} \in \mathcal{B}^{(q_i)}, \lambda_i \in \Pi\}$$

of $\mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ are linearly independent.

Proof. By Proposition 8.5 it is enough to prove that, for any fixed $q_1 > \dots > q_l$, the set

$$\{\pi_{ss}(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}) \dots \pi_{ss}(b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \mid b^{(q_i)} \in \mathcal{B}^{(q_i)}, \lambda_i \in \Pi\}$$

is linearly independent. But by the HN filtration again, we have (up to a scalar)

$$\begin{aligned} & \pi_{ss}(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)}) \dots \pi_{ss}(b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)}) \\ &= \sum_{\mathcal{H}_i \in \mathcal{C}_{q_i}} c_{\mathcal{H}_1}(\pi_{ss}(b^{(q_1)} \cdot T_{\lambda_1}^{(q_1)})) \dots c_{\mathcal{H}_l}(\pi_{ss}(b^{(q_l)} \cdot T_{\lambda_l}^{(q_l)})) [\mathcal{H}_1 \oplus \dots \oplus \mathcal{H}_l]. \end{aligned}$$

The corollary is now a consequence of Proposition 8.4. $\diamond$

We are finally ready to prove (8.1). Recall the notations of Section 8.2. Let us set $\hat{\mathbf{n}}_{>0} = \bigoplus_{q>0} \hat{\mathbf{n}}_q$ and let $\hat{\mathbf{n}}_0$ be the subalgebra generated by the root vectors of weight $\alpha_*, \tau(\alpha_*), \dots, \tau^{p-1}(\alpha_*)$. Similarly, let us denote by $\mathbf{U}_{\mathbf{p}, \Delta}^{(q)} \subset \mathbf{U}_{\mathbf{p}, \Delta}^{\geq 0}$ the vector space spanned by the elements $\{b^{(q)} \cdot T_{\lambda}^{(q)} \mid b \in \mathcal{B}^{(q)}, \lambda \in \Pi\}$, and by $\mathbf{U}_{\mathbf{p}, \Delta}^0$ the subalgebra generated by the line bundles $[\mathcal{O}], [\mathcal{O}(\vec{\omega})], \dots, [\mathcal{O}((p-1)\vec{\omega})]$.

By the PBW theorem we have, for any $\alpha \in \widehat{Q}$,

$$\begin{aligned} \dim \mathbf{U}^{\geq 0}(\widehat{\mathfrak{n}})[\alpha] &\leq \dim \mathbf{U}(\widehat{\mathfrak{n}}_0 \oplus \widehat{\mathfrak{n}}_{>0})[\alpha] \\ &= \sum_{\substack{\alpha=\alpha_0+\dots+\alpha_l \\ q_1>\dots>q_l>0}} \dim \mathbf{U}(\widehat{\mathfrak{n}}_0)[\alpha_0] \cdot \dim \mathbf{U}(\widehat{\mathfrak{n}}_{q_1})[\alpha_1] \cdots \dim \mathbf{U}(\widehat{\mathfrak{n}}_{q_l})[\alpha_l]. \end{aligned}$$

On the other hand, from Corollary 8.2 we deduce

$$\dim \mathbf{U}_{\mathbf{p},\Delta}^{\geq 0}[\alpha] \geq \sum_{\substack{\alpha=\alpha_0+\dots+\alpha_l \\ q_1>\dots>q_l>0}} \dim \mathbf{U}_{\mathbf{p},\Delta}^0[\alpha_0] \cdot \dim \mathbf{U}_{\mathbf{p},\Delta}^{(q_1)}[\alpha_1] \cdots \dim \mathbf{U}_{\mathbf{p},\Delta}^{(q_l)}[\alpha_l].$$

To conclude, observe that, by Corollary 8.1 and Section 7.4, we have for any β

$$\dim \mathbf{U}(\widehat{\mathfrak{n}}_q)[\beta] = \dim \mathbf{U}(\widehat{\mathfrak{t}})[\beta_\infty] = \dim \mathbf{U}_{\mathbf{p},\Delta}^{(\infty)}[\beta_\infty],$$

where β_∞ corresponds to β under the identification $\widehat{\Delta}_q \simeq \widehat{\Delta}_\infty$, and that by Lemma 8.3 and Proposition 8.4 we have

$$\dim \mathbf{U}_{\mathbf{p},\Delta}^{(\infty)}[\beta_\infty] = \dim \mathbf{U}_{\mathbf{p},\Delta}^{(q)}[\beta].$$

9. CONJECTURES

9.1. We first recall the Kac conjectures for representations of quivers. Let $Q = (I, \Omega)$ be an arbitrary quiver with no edge loops. Let $\mathfrak{g}$ be the Kac-Moody algebra associated to Q , let $\mathfrak{h} \subset \mathfrak{g}$ be a Cartan subalgebra and let $\Delta^+ \subset \mathfrak{h}^*$ denote the set of positive roots. We view the dimension $\mathbf{dim} M \in \mathbb{N}^I$ of some object $M \in \text{Rep}_k(Q)$ as an element of $\mathfrak{h}^*$ via the map $\mathbb{N}^I \rightarrow \mathfrak{h}^*$, $(n_i)_i \mapsto \sum_i n_i \alpha_i$.

Theorem 9.1 (Kac). *The following hold :*

- i) *There exists an indecomposable object in $\text{Rep}_k(Q)$ of dimension α if and only if $\alpha \in \Delta^+$. Moreover, the set of such indecomposables forms a $1 - \frac{(\alpha, \alpha)}{2}$ -parameter family with a unique component of maximal dimension.*
- ii) *For every root α there exists a polynomial $P_\alpha(v) \in \mathbb{Z}[v]$ with leading term $v^{1 - \frac{(\alpha, \alpha)}{2}}$ such that the number of absolutely indecomposable representations of dimension α over the finite field $\mathbb{F}_q$ is $P_\alpha(q)$.*

Conjecture (Kac). *We have $P_\alpha(v) \in \mathbb{N}[v]$ and $P_\alpha(0) = \dim \mathfrak{g}_\alpha$.*

This conjecture is trivial for finite type Dynkin quivers, proved for affine quivers and remains open in all other cases (see however [CB-VdB] for a recent important progress).

9.2. The results of this paper naturally lead us to the following variant of the above conjectures. Let $\mathbb{X}_{\mathbf{p},\Delta}$ be a weighted projective line and let $\mathfrak{L}\mathfrak{g}$ be the corresponding loop Kac-Moody algebra. Recall the identification $K_0(\text{Coh}(\mathbb{X}_{\mathbf{p},\Delta})) \simeq \mathfrak{h}_{\mathfrak{L}\mathfrak{g}}^*$ of Proposition 5.1.

Conjecture. *The following hold :*

- i) *There exists an indecomposable coherent sheaf $\mathcal{F} \in \text{Coh}(\mathbb{X}_{\mathbf{p},\Delta})$ if and only if $[\mathcal{F}] \in \mathfrak{h}_{\mathfrak{L}\mathfrak{g}}^*$ is a positive root. Moreover, the set of such indecomposables forms a $1 - \frac{(\alpha, \alpha)}{2}$ -parameter family with a unique component of maximal dimension.*

- ii) For every root α there exists a polynomial $P_\alpha(v) \in \mathbb{N}[v]$ with leading term $v^{1-\frac{(\alpha, \alpha)}{2}}$ such that the number of absolutely indecomposable coherent sheaves of dimension α over the finite field $\mathbb{F}_q$ is $P_\alpha(q)$, and we have $P_\alpha(0) = \dim \mathfrak{Lg}_\alpha$.

Conjecture i) was first made by Crawley-Boevey in relation to the Deligne-Simpson problem ([CB]). It is easy to check (using the descriptions of $\text{Coh}(\mathbb{X}_{\mathbf{p}, \underline{\lambda}})$ in Sections 7 and 8) that i) and ii) are true when $\mathbb{X}_{\mathbf{p}, \underline{\lambda}}$ is of genus $g \leq 1$.

10. APPENDIX

In this appendix we give the details of the calculations of Lemma 6.4. The computations of Section 6.9 are conducted in an exactly similar fashion and we leave them to the reader. For simplicity we drop the index (s) throughout in the notation (as in p_s for instance).

Proof of Lemma 6.4. Suppose that (6.3) holds for $r_1 = 1$ and arbitrary r_2 . Set $\tilde{T}_r = \frac{r}{[r]} T_r$. Then

$$[S_{1, r_2}] = [[S_1], \tilde{T}_{r_2}] = (-1)^{r_2} ad^{r_2} \tilde{T}_{r_2}([S_1]).$$

Since $[T_r, T_s] = 0$ for all r, s we deduce

$$\begin{aligned} [[S_{1, r_2}], \tilde{T}_{r_1}] &= (-1)^{r_2} [ad^{r_2} \tilde{T}_1([S_1]), \tilde{T}_{r_1}] \\ &= (-1)^{r_2} ad^{r_2} \tilde{T}_1([S_1], \tilde{T}_{r_1}) \\ &= (-1)^{r_2} ad^{r_2} \tilde{T}_1([S_{1, r_1}]) \\ &= S_{1, r_1 + r_2}. \end{aligned}$$

Thus it is enough to prove (6.3) for $r_2 = 1$. Let us introduce some notations. We label the boxes of a partition λ according to their height. For instance, the partition (32^21) is labelled

3			
2	2	2	
1	1	1	1

Let us write $\lambda \subset \nu$ if ν is obtained from λ by addition of a single box, and we write $\nu \setminus \lambda = \boxed{i}$ if the corresponding box is labelled by i . We also denote by $l(\lambda)_{\geq i}$ (resp. $l(\lambda)_{> i}$) the number of parts λ_j of $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$ with value $\lambda_j \geq i$ (resp. $\lambda_j > i$). Finally, if $\nu \setminus \lambda = \boxed{i}$ we set

$$p(\nu, \lambda) = \frac{v^{2l(\lambda)_{\geq i}} - v^{2l(\lambda)_{> i}}}{v^2 - 1}.$$

Lemma 10.1. *We have*

$$\tilde{T}_1 \tilde{T}_r = \sum_{\lambda, |\lambda|=r} n(l(\lambda) - 1) \sum_{\nu \supset \lambda} p(\nu, \lambda) [\nu].$$

Proof. It is enough to prove that

$$[[\lambda]] \cdot [[1]] = \sum_{\nu \supset \lambda} p(\nu, \lambda) [\nu].$$

By Section 4.2, we can assume $p = 1$. Let $(M)\nu, x$ be a representation of $\mathcal{C}_1$ of type ν . The space $N = \text{Ker } x$ has a natural filtration $N = N_0 \supset N_1 \supset \dots$ where

$N_i = N \cap \text{Im } x^i$. We have $\dim N_i = l(\nu)_{>i}$. Let $\lambda \subset \nu$ and let $L \subset N$ be a line. Then M_ν/L is of type λ if and only if $L \subset N_{i-1}$ but $L \not\subset N_i$ where $\nu \setminus \lambda = \boxed{i}$. Note that the number of such lines is $p(\nu, \lambda)$. $\diamond$

A similar computation shows that

$$\begin{aligned} [[\nu]] \cdot [S_1] &= v^{2l(\lambda)} [[\nu] \oplus S_1], \\ [S_1] \cdot [[\nu]] &= [[\nu] \oplus S_1] + \sum_{\gamma \triangleleft \nu} [[\nu \setminus \gamma] \oplus [1; p\gamma + 1]]. \end{aligned}$$

Hence,

$$(10.1) \quad \tilde{T}_1 \tilde{T}_r S_1 = \sum_{\lambda} n(l(\lambda) - 1) \sum_{\nu \supset \lambda} p(\nu, \lambda) v^{2l(\nu)} [[\nu] \oplus S_1],$$

$$(10.2) \quad \begin{aligned} S_1 \tilde{T}_1 \tilde{T}_r &= \sum_{\lambda} n(l(\lambda) - 1) \sum_{\nu \supset \lambda} p(\nu, \lambda) [[\nu] \oplus S_1] \\ &+ \sum_{\lambda} n(l(\lambda) - 1) \sum_{\nu \supset \lambda} p(\nu, \lambda) \sum_{\gamma \triangleleft \nu} [[\nu \setminus \gamma] \oplus [1; p\gamma + 1]]. \end{aligned}$$

A reasoning close to that of Lemma 10.1 shows that

$$\begin{aligned} [[\lambda] \oplus S_1] \cdot T_1 &= \sum_{\nu \supset \lambda} p(\nu, \lambda) [[\lambda] \oplus S_1] + v^{2l(\lambda) > 1} [[\lambda] \oplus [1; p + 1]], \\ [[1] \oplus S_1] \cdot [[\lambda]] &= \sum_{\nu \supset \lambda} p(\nu, \lambda) [[\nu] \oplus S_1] \\ &+ \sum_{\gamma \triangleleft \lambda} \sum_{\sigma \supset \lambda \setminus \gamma} p(\sigma, \lambda \setminus \gamma) [[\sigma] \oplus [1; p\gamma + 1]], \end{aligned}$$

from which one deduces

$$(10.3) \quad \begin{aligned} \tilde{T}_r S_1 \tilde{T}_1 &= \sum_{\lambda} n(l(\lambda) - 1) \sum_{\nu \supset \lambda} v^{2l(\lambda)} p(\nu, \lambda) [[\nu] \oplus S_1] \\ &+ \sum_{\lambda} n(l(\lambda) - 1) v^{2(l(\lambda) + l(\lambda) > 1)} [[\lambda] \oplus [1; p + 1]], \end{aligned}$$

$$(10.4) \quad \begin{aligned} \tilde{T}_1 S_1 \tilde{T}_r &= \sum_{\lambda} v^2 n(l(\lambda) - 1) \sum_{\nu \supset \lambda} p(\nu, \lambda) [[\nu] \oplus S_1] \\ &+ \sum_{\lambda} v^2 n(l(\lambda) - 1) \sum_{\gamma \triangleleft \lambda} \sum_{\sigma \supset \lambda \setminus \gamma} p(\sigma, \lambda \setminus \gamma) [[\sigma] \oplus [1; p\gamma + 1]]. \end{aligned}$$

We are now in position to prove Lemma 6.4. We have

$$(10.5) \quad [S_{1,r}, \tilde{T}_1] = S_1 \tilde{T}_r \tilde{T}_1 + \tilde{T}_r \tilde{T}_1 S_1 - \tilde{T}_r S_1 \tilde{T}_1 - \tilde{T}_1 S_1 \tilde{T}_r.$$

The coefficient c_ν of $[[\nu] \oplus S_1]$ in (10.5) is

$$\begin{aligned} c_\nu &= (1 - v^2) \sum_{\lambda \subset \nu} n(l(\lambda) - 1) p(\nu, \lambda) + \sum_{\lambda \subset \nu} n(l(\lambda) - 1) p(\nu, \lambda) (v^{2l(\nu)} - v^{2l(\lambda)}) \\ &= (1 - v^2) \sum_{\substack{\lambda \subset \nu \\ \nu \setminus \lambda \neq 1}} n(l(\nu) - 1) p(\nu, \lambda) \\ (10.6) \quad &+ \sum_{\substack{\lambda \subset \nu \\ \nu \setminus \lambda = 1}} n(l(\nu) - 2) (1 - v^2) (1 - v^{2(l(\nu) - 1)}) p(\nu, \lambda) \\ &= (1 - v^2) n(l(\nu) - 1) \sum_{\lambda \subset \nu} p(\nu, \lambda) \\ &= n(l(\nu)) \end{aligned}$$

since

$$\sum_{\lambda \subset \nu} p(\nu, \lambda) = \frac{v^{2l(\nu)} - 1}{v^2 - 1}.$$

Similarly, the coefficient $c_{\nu, \gamma}$ of $[\nu] \oplus [1; p\gamma + 1]$ in (10.5) is, if $\gamma > 1$,

$$(10.7) \quad c_{\nu, \gamma} = \sum_{\lambda \subset \nu \cup \{\gamma\}} n(l(\lambda) - 1) p(\nu \cup \{\gamma\}, \lambda) - v^2 \sum_{\sigma \subset \nu} n(l(\sigma)) p(\nu, \sigma).$$

Observe that

$$p(\nu, \sigma) = \begin{cases} p(\nu \cup \{\gamma\}, \sigma \cup \{\gamma\}) & \text{if } \nu \setminus \sigma > \boxed{\gamma}, \\ v^{-2} p(\nu \cup \{\gamma\}, \sigma \cup \{\gamma\}) & \text{if } \nu \setminus \sigma < \boxed{\gamma}, \\ p(\nu \cup \{\gamma\}, \sigma \cup \{\gamma\}) - v^{2(l(\nu) \geq \gamma)} & \text{if } \nu \setminus \sigma = \boxed{\gamma} \end{cases}$$

substituting in (10.7) yields

$$(10.8) \quad \begin{aligned} c_{\nu, \gamma} &= n(l(\nu)) \left\{ (1 - v^2) \sum_{\substack{\sigma \subset \nu \\ \nu \setminus \sigma \geq \gamma}} p(\nu \cup \{\gamma\}, \sigma \cup \{\gamma\}) + v^{2(l(\nu) \geq \gamma + 1)} \right\} \\ &= n(l(\nu)) \left\{ (1 - v^2) \frac{v^{2(l(\nu) \geq \gamma + 1)} - 1}{v^2 - 1} + v^{2(l(\nu) \geq \gamma + 1)} \right\} \\ &= n(l(\nu)). \end{aligned}$$

Similar computation shows that (10.8) holds for $\gamma = 1$ also. Equations (10.6) and (10.8) together prove Lemma 6.4. $\diamond$

REFERENCES

- [BK] Baumann, P., Kassel, C., *The Hall algebra of the category of coherent sheaves on the projective line.*, J. Reine Angew. Math. **533** 207-233, (2001).
- [Be] Beck J., *Braid group action and quantum affine algebras.*, Comm. Math. Phys. **165** no. 3, 555-568, (1994).
- [Bo] Bondal A.I., *Representations of associative algebras and coherent sheaves*, Izv. Akad. Nauk. SSSR Ser. Math. **53**, 25-44, (1989).
- [CB] Crawley-Boevey B., *On the Deligne-Simpson problem and weighted projective lines*, in preparation (Talk at ICRA X 2002).
- [CB-VdB] Crawley-Boevey B., Van den Bergh M., *Absolutely indecomposable representations and Kac-Moody algebras*, preprint RA/0106009.
- [E] Enriquez B., *PBW and duality theorems for quantum groups and quantum current algebras*, preprint QA/9904113.
- [GL] Geigle W., Lenzing H., *A class of weighted projective curves arising in the representation theory of finite-dimensional algebras*, Singularities, representations of algebras and vector bundles, Springer LNM **1273**, 265-297, (1987).
- [Hal] Hall P., *The algebra of partitions*, Proc. 4th Canadian Math. Congress, 147-159, (1959).
- [Hap1] Happel, D. *Triangulated categories in the representation theory of finite dimensional algebras*, London Math. Soc. Lect. Notes **119** (1988).
- [Hap2] Happel, D., *A characterization of hereditary categories with tilting object*, Invent. Math. **144** no. 2, 381-398, (2001).
- [Kac] Kac V., *Infinite dimensional Lie algebras*, Cambridge Univ. Press, (1990).
- [Kap] Kapranov, M., *Eisenstein series and quantum affine algebras*, Algebraic geometry 7, J. Math. Sci. **84** no. 5, 1311-1360, (1997).
- [La] Lamotke, K., *Regular solids and isolated singularities*. Adv. Lect. in Math., Braunschweig, (1986).
- [LT] Leclerc B., Thibon J-Y., *Littlewood-Richardson coefficients and Kazhdan-Lusztig polynomials.*, Combinatorial methods in representation theory (Kyoto, 1998), 155-220, Adv. Stud. Pure Math. **28**.
- [Le] Lenzing H., *Curve singularities arising from the representation theory of tame hereditary algebras*, Representation theory, I (Ottawa, Ont., 1984), Lecture Notes in Math. **1177**, 199-231.

- [LM] Lenzing H., Meltzer H., *Sheaves on a weighted projective line of genus one, and representations of a tubular algebra*, Representations of algebras (Ottawa, ON, 1992), 313-337, CMS Conf. Proc., **14**.
- [LP] Lin Y., Peng L., *2-Extended affine Lie algebras and tubular algebras*, to appear in Adv. Math..
- [Lu] Lusztig G., *Introduction to quantum groups*, Progress in Mathematics, **110**. Birkhauser, (1993).
- [MRY] Moody R., Rao S., Yokonuma Y., *Toroidal Lie algebras and vertex representations*, Geom. Dedicat. **35**, 283-307 (1990).
- [N] Nakajima, H., *Quiver varieties and Kac-Moody algebras.*, Duke Math. J. **91**, no. 3, 515-560, (1998).
- [M] Macdonald I., *Symmetric functions and Hall polynomials*, Oxford press (1979).
- [PX] Peng, L., Xiao, J., *Triangulated categories and Kac-Moody algebras.*, Invent. Math. **140**, no. 3, 563-603, (2000).
- [Re] Reineke M., *The Harder-Narasimhan system in quantum groups and cohomology of quiver moduli*, math.QA/0204059.
- [Ri1] Ringel C., *Hall algebras and quantum groups*, Invent. Math. **101** no. 3, 583-591, (1990).
- [Ri2] Ringel C., *Hall algebras*, Topics in algebra, Part 1 (Warsaw 1988), PWN, Warsaw, 433-447, (1990).
- [Ri3] Ringel C., *Tame algebras and integral quadratic forms*, LNM **1099**, (1984).
- [S] Schiffmann O., *The Hall algebra of a cyclic quiver and canonical bases of Fock spaces.*, Internat. Math. Res. Notices, no. **8**, 413-440, (2000).

Olivier Schiffmann, schiffma@dma.ens.fr,
 Yale University, 10 Hillhouse Avenue, 06520-8283 New Haven, CT, USA, and
 DMA, École Normale Supérieure, 45 rue d'Ulm, 75230 Paris Cedex 05-FRANCE